Natural Resources for Human Health
eISSN: 2583-1194

Original Research

Received 09 July 2026
Revised 24 Aug 2026
Accepted 07 Sep 2026

Natural Resources for Human Health 2026, Vol. 6 (19s): 596–615

KEYWORDS:
Radiology; Pharmacy; Medical Nurse; Medical Secretary; Health Information; Diagnostic Imaging; Patient-Centered Care; Imaging Safety; Radiation Protection; Contrast Media; Medication Safety; Patient Preparation; Magnetic Resonance Imaging Safety; Computed Tomography; Medication Reconciliation; Diagnostic Accuracy; Electronic Health Records; Radiology Information Systems; Picture Archiving and Communication Systems; Clinical Communication; Patient Education; Diagnostic Errors; Healthcare Coordination; Quality Improvement; Continuity of Care.

Radiology, Pharmacy, Medical Nurse, Medical Secretary, and Health Information Collaboration for Comprehensive Diagnostic Imaging Safety and Patient-Centered Care

Obeid Mohammed Laslom[1], Sara Olyan Al-Rashidi[2], Hamad Khamsan Ali Althaiban[3], Lamees Tariq Alghamdi[4], Rania Mohammed Ghaleb[5], Ibtihal Hawash Alaradi[6], Sarah Abdullah Saud Alotaibi[7], Aqelah Hayder Alhashem[8], Abdullah Mohammad Mufrah Asere[9], Amal Saleh Algarni[10], Amal Mohammad Alqahtani[11], Ziyad Helal Alotaibi[12], Duha Nabeel Bukhari[13], Fahad Shabab Alghamdi[14], Mohammed Nayidh Juraybia Alresheedi[15], Satem Abdullah A Aloraini[16]

[1]Forensic Medical Services Center Ministry Of Health Kingdom Of Saudi Arabia
[2]Qassim University Medical City Ministry Of Education Kingdom Of Saudi Arabia
[3]Hada Prison Ministry Of Interior Kingdom Of Saudi Arabia
[4]10,11-King Abdulaziz Air Base Military Hospital Ministry Of Defense Kingdom Of Saudi Arabia
[5]Armed Forces Hospital Ministry Of Defense Kingdom Of Saudi Arabia
[6,7,8]King Fahd Armed Forces Hospital Ministry Of Defense Kingdom Of Saudi Arabia
[9]Abha Prison Health Center Ministry Of Interior Kingdom Of Saudi Arabia
[12]Alkharj Military Hospital Ministry Of Defense Kingdom Of Saudi Arabia
[13]National Guard Hospital Ministry Of National Guard Kingdom Of Saudi Arabia
[14]King Abdulaziz University Hospital Ministry Of Education Kingdom Of Saudi Arabia
[15,16]Armed Forces Hospital In Al-Qassim Ministry Of Defense Kingdom Of Saudi Arabia

ABSTRACT: Diagnostic imaging has become an indispensable component of contemporary healthcare, supporting disease detection, clinical decision-making, treatment planning, and the evaluation of therapeutic outcomes. The increasing use of radiography, computed tomography, magnetic resonance imaging, ultrasonography, and other imaging modalities has improved diagnostic capabilities while introducing additional responsibilities concerning patient preparation, medication management, procedural safety, clinical communication, and continuity of care. Although technological advancements have enhanced imaging accuracy and efficiency, the safety and overall experience of patients depend on much more than the performance of imaging equipment. Delays in scheduling, incomplete medical histories, inappropriate medication management, inadequate screening for contraindications, failures in communicating critical findings, and fragmented documentation can compromise diagnostic quality and expose patients to preventable harm. These challenges are particularly important among older adults, children, pregnant patients, individuals with renal impairment, patients

Natural Resources for Human Health
eISSN: 2583-1194

receiving complex pharmacological treatment, and those requiring repeated or specialized imaging examinations.

Comprehensive diagnostic imaging safety requires coordinated participation from several healthcare disciplines whose responsibilities extend across the entire imaging pathway. Radiology professionals contribute through appropriate examination selection, protocol optimization, radiation protection, contrast administration oversight, image acquisition, and accurate interpretation. Pharmacy professionals support medication reconciliation, assessment of relevant drug interactions, contrast-related medication considerations, and the safe use of sedatives, analgesics, and emergency medicines when clinically indicated. Medical Nurses provide patient assessment, preparation, education, clinical monitoring, recognition of adverse reactions, and post-procedural care. Medical Secretaries contribute through referral processing, appointment coordination, verification of administrative information, communication of preparation instructions, and organization of follow-up arrangements. Health Information professionals maintain the accuracy, availability, confidentiality, and interoperability of clinical records, imaging documentation, and diagnostic reports.

The effectiveness of this collaborative approach depends on clearly defined responsibilities, timely information exchange, standardized safety procedures, and attention to individual patient needs. Patient-centered diagnostic imaging requires consideration of physical comfort, psychological well-being, communication preferences, informed decision-making, accessibility, cultural sensitivity, and the patient's ability to understand and complete recommended investigations. Particular attention must be given to the safe use of contrast media, prevention of medication-related complications, radiation dose optimization, MRI safety screening, management of vulnerable patients, and reliable communication of urgent imaging findings.

This review examines the complementary contributions of Radiology, Pharmacy, Medical Nurse, Medical Secretary, and Health Information departments to diagnostic imaging safety and patient-centered care. It discusses the clinical and operational risks associated with fragmented imaging pathways and explores how coordinated assessment, preparation, monitoring, documentation, and follow-up can strengthen diagnostic reliability. The article also considers the role of digital health systems, quality improvement, staff education, and institutional governance in supporting safer imaging services. Integrating these professional functions into a continuous care pathway offers a practical means of reducing preventable complications, improving patient experiences, and ensuring that diagnostic information is translated into timely and appropriate clinical decisions.

1. INTRODUCTION

Diagnostic imaging occupies a central position in modern clinical practice. Physicians rely on imaging examinations to investigate symptoms, confirm or exclude suspected diseases, assess anatomical and functional abnormalities, guide therapeutic interventions, and monitor changes in a patient's condition over time. Developments in computed tomography (CT), magnetic resonance imaging (MRI), ultrasonography, digital radiography, mammography, and image-guided procedures have expanded the ability of healthcare professionals to identify abnormalities that might previously have remained undetected until more advanced stages of disease. In emergency medicine, imaging can provide information necessary for immediate decisions involving trauma, stroke, internal bleeding, and other potentially life-threatening conditions. In chronic disease management, it contributes to diagnosis, surveillance, treatment evaluation, and long-term clinical planning.

Despite these benefits, diagnostic imaging is not an isolated technical activity. Every examination forms part of a wider clinical process that begins when an investigation is considered and continues through referral, scheduling, patient assessment, preparation, image acquisition, interpretation, communication of findings, and appropriate follow-up. Errors or omissions at any stage can diminish the value of an otherwise technically successful examination. A correctly performed CT scan, for example, may still fail to improve patient outcomes if the examination was inappropriate, essential clinical information was unavailable, a significant finding was not communicated promptly, or the patient did not receive the recommended follow-up care.

The quality of diagnostic imaging must therefore be evaluated in relation to both the accuracy of the examination and the reliability of the care pathway surrounding it. Patient safety includes protection from unnecessary radiation exposure, prevention of contrast-related complications, recognition of medication-related risks, appropriate screening before MRI examinations, safe patient positioning and transfers, infection prevention, and timely management of unexpected clinical deterioration. Equally important are the administrative and informational processes that determine whether patients receive the correct examination, arrive adequately prepared, and obtain appropriate clinical attention after the results become available.

These responsibilities cannot be fulfilled effectively by Radiology alone. The contributions of Pharmacy, Medical Nurse, Medical Secretary, and Health Information departments are essential to addressing risks that arise before, during, and after diagnostic procedures. Their collaboration becomes particularly important when patients have multiple chronic illnesses, complicated medication regimens, limited mobility, communication difficulties, or conditions requiring additional clinical supervision.

1.1 Evolution of Diagnostic Imaging and Its Implications for Patient Safety

The development of modern imaging technology has substantially changed the way healthcare professionals investigate disease. Conventional radiography remains important for evaluating fractures, pulmonary abnormalities, and numerous other clinical conditions, while CT provides rapid cross-sectional imaging that is particularly valuable in emergency and complex diagnostic situations. MRI offers detailed soft-tissue assessment without ionizing radiation, and ultrasonography provides real-time examination for many abdominal, vascular, obstetric, and musculoskeletal indications.

These modalities differ in their clinical applications, safety considerations, preparation requirements, and potential complications. Consequently, an examination that is appropriate for one patient may be unsuitable for another.

CT examinations involving ionizing radiation require justification and optimization of exposure, with particular attention to children and patients undergoing repeated investigations. MRI introduces different concerns, including the presence of certain implants, metallic foreign bodies, and equipment that may be unsafe within the magnetic environment. Contrast-enhanced studies require assessment of previous reactions and relevant patient-specific risk factors. Some procedures involve intravenous access, sedation, analgesia, or post-procedural observation.

The growing complexity of diagnostic imaging has increased the importance of reliable clinical information and coordinated professional judgment. Technological sophistication does not eliminate human or organizational error. In some circumstances, it makes accurate preparation and communication even more important because an examination may involve multiple decisions and handovers within a relatively short period.

A comprehensive approach to imaging safety must account for these differences without treating every imaging examination as having the same level of risk.

1.2 Diagnostic Imaging as a Continuum of Patient Care

The imaging pathway begins before the patient enters the Radiology department. A clinician must determine whether an examination is clinically indicated, select an appropriate modality, and provide sufficient information for the imaging team to understand the diagnostic question.

The referral then enters an administrative and clinical workflow involving appointment allocation, examination-specific preparation, verification of relevant patient information, and assessment of possible safety concerns.

Medical Secretaries are often responsible for ensuring that referral documents are complete and that appointments are organized according to established scheduling and prioritization procedures. They may communicate standardized preparation

instructions and direct clinical questions to the appropriate professionals. They should not independently determine clinical suitability or interpret contraindications.

Medical Nurses assess clinical needs, identify relevant risks, prepare patients, and provide support during procedures that require nursing involvement. Pharmacists assist when medication history, treatment modification, drug interactions, or pharmacological monitoring may affect the examination.

Health Information professionals support the availability and integrity of the clinical records required throughout the process.

Radiology professionals then perform and interpret the examination, with responsibility for ensuring that the selected protocol addresses the clinical question while minimizing avoidable risk.

The process does not end when images have been acquired. Findings must be reported, communicated, incorporated into the patient's medical record, and acted upon by the responsible clinical team. Incomplete follow-up can undermine the entire diagnostic process, even when image acquisition and interpretation have been technically satisfactory.

1.3 Patient-Centered Care in Diagnostic Imaging

Patient-centered care recognizes that individuals undergoing imaging examinations are not simply recipients of diagnostic procedures. They may be experiencing pain, uncertainty, physical limitations, emotional distress, or fear about the possibility of serious disease.

Some patients are unfamiliar with imaging technology and may be concerned about radiation exposure, contrast injections, enclosed scanning environments, or the potential meaning of their results. Others may have experienced previous difficult examinations or may struggle to follow preparation instructions because of language barriers, sensory impairment, limited health literacy, or cognitive difficulties.

A patient-centered imaging service should provide clear information about the purpose of the examination, expected procedure, relevant risks, preparation requirements, and arrangements for receiving results. Communication should be adapted to the patient's needs rather than relying exclusively on standard written instructions.

Medical Nurses can address clinical concerns and provide reassurance before and during procedures. Medical Secretaries contribute by ensuring that appointment information and authorized preparation instructions are communicated clearly and consistently. Radiology professionals explain procedural requirements and respond to questions within their responsibilities.

Pharmacy professionals may provide individualized counseling when medication management affects preparation or recovery. Health Information systems support access to relevant patient preferences, previous examinations, and documented clinical information.

Patient-centered care also requires respect for privacy, dignity, informed consent where applicable, and the patient's right to participate in decisions concerning their care.

These considerations are particularly important during examinations involving intimate positioning, prolonged immobility, intravenous access, or sedation.

1.4 Patient Safety Challenges Across the Imaging Pathway

Preventable harm in diagnostic imaging can arise from several interconnected sources. Some relate directly to imaging technology, while others originate from communication failures, inadequate preparation, or incomplete clinical information.

An examination may be performed on the wrong patient or anatomical site if identification and verification procedures are insufficient. An incomplete referral may lead to selection of an unsuitable imaging protocol. Failure to recognize a relevant previous contrast reaction may expose a patient to avoidable risk. Missing information about an implanted medical device can create a serious MRI safety concern.

Medication-related problems may arise when examination preparation affects usual treatment, when sedation is required, or when the patient's clinical condition alters the safety of prescribed medicines.

Patients with limited mobility may experience falls or injuries during transfer or positioning. Individuals with significant respiratory or cardiovascular disease may require closer monitoring during certain procedures.

Operational problems also affect safety. Delayed appointments, inadequate communication of preparation instructions, unavailable previous imaging studies, and failures to communicate urgent findings can prolong diagnosis or delay treatment.

These risks demonstrate why imaging safety should be understood as a shared clinical and organizational responsibility.

1.5 The Role of Radiology in Diagnostic Quality and Procedural Safety

Radiology is responsible for the clinical and technical integrity of diagnostic imaging. Its responsibilities include reviewing examination appropriateness, selecting suitable protocols, optimizing image quality, managing modality-specific safety requirements, and ensuring accurate interpretation.

Radiologists and radiographers or radiologic technologists have distinct professional responsibilities, which vary according to local regulation and institutional policy. Their work must be coordinated to ensure that imaging examinations answer the clinical question without exposing patients to unnecessary risk.

Radiation protection is particularly important in examinations involving ionizing radiation. Imaging protocols should be justified and optimized according to the patient's clinical needs, age, body size, and the diagnostic information required.

In MRI, appropriate screening is necessary to identify implants, devices, and other potential hazards before entering controlled areas.

For contrast-enhanced examinations, Radiology must ensure that the indication, contrast protocol, relevant clinical history, and available emergency arrangements are appropriate.

The department also has a critical responsibility to communicate urgent or unexpected findings through established escalation pathways.

A radiology report is clinically valuable only when its findings are accessible, understandable, and appropriately incorporated into patient management.

1.6 Pharmacy and Medication Safety in Diagnostic Imaging

Pharmacy contributes to diagnostic imaging safety by addressing pharmacological issues that may influence examination preparation, procedural management, and recovery.

Many patients undergoing imaging receive medicines for chronic conditions, including diabetes, cardiovascular disease, neurological disorders, and chronic pain. Some may require individualized medication instructions because of fasting, sedation, altered oral intake, renal impairment, or planned interventional procedures.

Medication reconciliation can identify discrepancies between prescribed treatment and medicines actually being taken, including over-the-counter products and other substances relevant to clinical assessment.

Pharmacists can help evaluate potential drug interactions, advise on medication-related precautions, and support the safe use of sedatives, analgesics, and emergency medicines where indicated.

Contrast-related medication management requires particular care. Decisions should reflect the contrast agent, the examination, the patient's renal function and other clinical characteristics, and current institutional guidance. Broad instructions to stop commonly used medicines without individualized assessment may create unnecessary clinical risk.

Pharmacy also supports the availability and safe storage of medications required for managing acute adverse reactions.

Its contribution is most effective when medication recommendations are communicated before the patient arrives for the examination, allowing necessary changes to be implemented safely and without avoidable delay.

1.7 Medical Nurse Responsibilities in Patient Preparation and Monitoring

Medical Nurses provide an important connection between diagnostic procedures and the patient's broader clinical condition.

Before an examination, nursing assessment may identify symptoms, relevant allergies, previous adverse reactions, mobility limitations, anxiety, communication difficulties, or conditions requiring additional observation.

For patients undergoing contrast-enhanced procedures, sedation, or interventions, Nurses may assist with preparation, intravenous access, administration of authorized medications, and monitoring according to professional scope and institutional policy.

During procedures, Nursing involvement is particularly important when patients are clinically unstable, unable to communicate effectively, or at increased risk of complications.

Nurses also contribute to safe patient transfers, infection prevention, assessment of discomfort, and recognition of unexpected changes in clinical status.

Following an examination, they may monitor recovery, identify delayed symptoms requiring assessment, provide discharge instructions, and communicate relevant observations to the treating team.

The continuity provided by Nursing helps ensure that diagnostic imaging remains connected to the patient's ongoing care rather than becoming an isolated departmental activity.

1.8 Medical Secretary and the Organization of Safe Imaging Services

Administrative coordination is sometimes overlooked in discussions of patient safety, yet errors in appointment management and communication can have significant clinical consequences.

Medical Secretaries contribute to safe imaging services by processing referrals, verifying administrative details, coordinating appointments, communicating approved preparation instructions, and ensuring that relevant documentation reaches the appropriate professionals.

For examinations requiring specific preparation, such as fasting or completion of preliminary investigations, clear administrative communication helps reduce cancellations, unnecessary repeat visits, and procedural delays.

Medical Secretaries may also identify incomplete referrals and direct them to authorized clinical staff for clarification.

Their involvement becomes particularly important when patients require urgent appointments, multiple examinations, interpreter services, transport arrangements, or coordinated follow-up.

They should not be expected to make clinical decisions about medication discontinuation, contrast suitability, pregnancy-related imaging risks, or emergency prioritization without an approved clinical pathway.

Clearly defined boundaries allow administrative personnel to contribute effectively while ensuring that clinical decisions remain with appropriately qualified professionals.

1.9 Health Information and the Integrity of Diagnostic Data

Diagnostic imaging depends on accurate and accessible clinical information.

Health Information professionals support the management of electronic health records, radiology information systems, picture archiving and communication systems, diagnostic reports, and associated clinical documentation.

Their responsibilities may include data quality, record integrity, patient identification processes, confidentiality, information governance, and the coordination of information exchange between clinical systems.

Errors in demographic information, duplicate patient records, missing reports, or incorrectly linked images can create serious risks.

A reliable information system should allow authorized professionals to review relevant clinical history, previous imaging examinations, allergies, and available investigation results when these are needed for safe care.

Health Information professionals also contribute to standardized documentation, report tracking, audit processes, and secure access to sensitive patient information.

The increasing use of digital imaging and electronic communication makes this role especially important, as diagnostic safety depends not only on generating accurate information but also on ensuring that it reaches the correct clinical team.

1.10 Importance of Coordinated Communication Between Departments

Communication failures are a recurring concern across complex healthcare pathways.

An imaging appointment may be delayed because preparation instructions were incomplete. A procedure may be interrupted because a medication issue was not identified in advance. An urgent finding may remain unaddressed because the responsible clinician did not receive timely notification.

These problems often arise at the boundaries between departments rather than within a single professional activity.

A coordinated communication system should define which information must be shared, who is responsible for sharing it, how urgent concerns are escalated, and how completion of critical tasks is confirmed.

For example, a patient scheduled for contrast-enhanced imaging may require clinical screening by Radiology, medication review by Pharmacy when indicated, nursing assessment, administrative coordination of the appointment, and availability of accurate clinical records.

If these activities occur independently without reliable communication, important information may be overlooked or duplicated.

Structured handovers, standardized referral requirements, documented clinical decisions, and closed-loop communication of urgent findings can reduce these risks.

The purpose of collaboration is not to increase the number of professionals involved in every examination, but to ensure that the right expertise is available when it is needed.

1.11 Diagnostic Imaging Safety in Vulnerable Patient Populations

Certain patient groups require additional consideration because their clinical conditions or circumstances may increase the complexity of diagnostic imaging.

Older adults may have multiple chronic illnesses, reduced mobility, sensory impairment, cognitive difficulties, or extensive medication regimens. These factors can complicate preparation, positioning, communication, and post-procedural observation.

Children require developmentally appropriate communication, suitable imaging protocols, and particular attention to radiation dose optimization. In some circumstances, additional support may be necessary to reduce movement and obtain diagnostically useful images without unnecessary repeat exposure.

Pregnant patients require examination selection based on clinical necessity, the body region involved, and the relative benefits and risks of available imaging modalities.

Patients with kidney disease may need individualized assessment when certain contrast agents are considered. Those with implanted medical devices require careful MRI safety screening.

Individuals with disabilities, limited language proficiency, or low health literacy may need adapted instructions, interpreters, accessible facilities, or additional assistance during appointments.

Coordinated involvement of the five departments can help address these needs without introducing unnecessary barriers to diagnostic care.

1.12 Diagnostic Delays, Missed Appointments, and Continuity of Care

The consequences of delayed imaging extend beyond inconvenience.

When clinically necessary examinations are postponed, diagnosis and treatment may also be delayed. Repeated cancellations can increase patient anxiety, create additional travel and financial burdens, and contribute to loss of follow-up.

Some delays arise from limited imaging capacity, but others reflect incomplete referrals, unavailable clinical information, unclear preparation instructions, or ineffective appointment coordination.

Medical Secretaries can identify scheduling problems and support timely communication with patients and clinical departments.

Medical Nurses and Pharmacy professionals may help resolve preparation-related clinical concerns before the scheduled examination.

Radiology determines appropriate examination protocols and clinical prioritization, while Health Information systems support tracking of orders, completed studies, reports, and outstanding follow-up requirements.

Reliable continuity of care requires particular attention to patients who miss appointments or do not receive recommended follow-up after abnormal imaging findings.

A completed imaging examination should be regarded as one stage of the diagnostic process, not necessarily its endpoint.

1.13 Quality Improvement and Institutional Responsibility

Comprehensive imaging safety depends on organizational systems that support consistent professional practice.

Hospitals and diagnostic centers should establish clear procedures for patient identification, examination verification, radiation protection, MRI screening, contrast safety, medication management, incident reporting, and communication of urgent findings.

Quality improvement requires regular evaluation of these processes rather than assuming that the absence of reported incidents demonstrates adequate safety.

Relevant indicators may include examination cancellations related to inadequate preparation, patient identification discrepancies, contrast-related incidents, repeat imaging due to avoidable technical or administrative errors, delayed communication of critical findings, and incomplete follow-up.

Patient feedback can provide additional insight into communication, comfort, accessibility, waiting times, and the clarity of preparation instructions.

The participation of Radiology, Pharmacy, Medical Nurse, Medical Secretary, and Health Information departments in quality-improvement activities can help identify problems that would otherwise remain hidden within individual workflows.

Leadership support is essential for translating findings into changes in staffing, training, technology, communication procedures, and accountability.

1.14 Rationale and Objectives of the Present Review

The increasing complexity of diagnostic imaging has created a need to examine patient safety from a broader perspective than technical image quality alone.

Radiology remains central to examination selection, procedural safety, and diagnostic interpretation, but the effectiveness of imaging services also depends on medication management, clinical monitoring, administrative coordination, and reliable health information.

Fragmentation among these functions can create preventable risks, particularly for patients with complex medical needs or those undergoing repeated examinations.

The present review therefore explores how collaboration among Radiology, Pharmacy, Medical Nurse, Medical Secretary, and Health Information professionals can strengthen diagnostic imaging safety while promoting patient-centered care.

Its objectives are to examine the responsibilities of each department, identify common safety challenges across the imaging pathway, discuss opportunities for coordinated prevention of diagnostic and procedural errors, and consider practical approaches to improving communication, documentation, patient preparation, and follow-up.

The review also addresses the importance of individualized care for vulnerable populations, appropriate use of digital information systems, and continuous quality improvement.

By examining these interconnected responsibilities, the article seeks to provide a practical framework for safer and more responsive diagnostic imaging services, in which technical accuracy, clinical judgment, operational reliability, and patient experience are treated as complementary dimensions of healthcare quality.

2. ESTABLISHING AN INTEGRATED DIAGNOSTIC IMAGING CARE PATHWAY

Diagnostic imaging is most effective when it is treated as part of a continuous clinical pathway rather than an isolated technical service. The process begins with a clinical question and ends only when the imaging findings have been appropriately communicated, interpreted within the patient's broader condition, and incorporated into subsequent care. Between these stages, numerous decisions influence patient safety, including examination selection, appointment prioritization, medication management, patient preparation, procedural monitoring, and documentation.

The complexity of this pathway becomes particularly apparent when patients have multiple medical conditions or require specialized investigations. A patient referred for contrast-enhanced computed tomography, for example, may have diabetes, impaired renal function, a previous contrast reaction, and limited mobility. Although the radiological examination itself may take only a short time, safe completion requires consideration of clinical history, current medication, vascular access, mobility assistance, examination preparation, and arrangements for communicating results.

Radiology professionals determine the appropriate imaging technique and examination protocol. Pharmacy evaluates medication-related concerns and supports safe pharmacological management when needed. Medical Nurses assess patient readiness, identify clinical risks, and provide monitoring during procedures requiring nursing involvement. Medical Secretaries organize appointments, verify referral completeness, and communicate approved instructions. Health Information professionals maintain the availability and accuracy of the clinical records needed for these activities.

A coordinated pathway should define responsibility at each stage without creating unnecessary duplication. For routine examinations, most patients will not require direct involvement from every department. However, systems should ensure that specialist support is readily available when clinical complexity or identified risk makes it necessary.

The central objective is to provide the right examination, to the right patient, at the right time, with appropriate preparation and reliable follow-up.

3. CLINICAL APPROPRIATENESS AND JUSTIFICATION OF IMAGING EXAMINATIONS

The safety of diagnostic imaging begins with deciding whether an examination is necessary and whether the selected modality is appropriate for the clinical question. Technically accurate imaging may still represent poor care when the investigation offers little diagnostic benefit, duplicates recent examinations without justification, or exposes the patient to avoidable risks.

Imaging appropriateness requires consideration of the suspected diagnosis, clinical urgency, patient characteristics, previous investigations, and available alternatives. In some situations, ultrasonography or MRI may provide sufficient diagnostic information without ionizing radiation. In others, CT may offer substantial clinical benefit because of its speed, accessibility, or ability to identify urgent pathology.

Radiology professionals contribute by reviewing referral indications, recommending suitable protocols, and advising on alternative examinations when appropriate. Referring clinicians retain responsibility for the clinical question and broader treatment decisions.

Health Information systems can improve this process by making previous imaging studies and relevant clinical findings accessible. Reviewing earlier examinations may prevent unnecessary duplication and help determine whether additional imaging is likely to change patient management.

Medical Secretaries support the process by identifying incomplete referral documentation and directing it for clinical clarification. Their role is particularly important when missing information would otherwise lead to appointment cancellation or inappropriate scheduling.

Patient-centered imaging also requires explaining why an examination is recommended and what information it is expected to provide. This helps patients understand the purpose of the investigation and participate meaningfully in decisions about their care.

4. PATIENT IDENTIFICATION AND PREVENTION OF WRONG-PATIENT IMAGING

Patient identification errors represent a serious risk because they can affect every stage of diagnostic care, from referral and scheduling to image acquisition, reporting, and subsequent treatment.

Incorrect demographic information, duplicate medical records, similar patient names, and incomplete verification procedures may result in examinations being performed on the wrong individual or images being associated with an incorrect record.

A reliable identification process should use approved patient identifiers at appropriate points in the imaging pathway. Administrative verification during appointment registration should be followed by clinical and procedural confirmation before the examination.

Medical Secretaries contribute through accurate registration and correction of administrative discrepancies. Medical Nurses and Radiology personnel verify identity before procedures, medication administration, or other clinically significant actions.

Health Information professionals help maintain data integrity across electronic health records, radiology information systems, and picture archiving and communication systems.

Particular care is necessary for patients who cannot reliably confirm their identity because of unconsciousness, cognitive impairment, language barriers, or critical illness.

Identification procedures should remain consistent during emergencies while allowing clinically appropriate adaptations for urgent care.

5. COMPREHENSIVE PRE-IMAGING CLINICAL ASSESSMENT

Pre-imaging assessment allows healthcare professionals to identify factors that may influence examination safety, preparation, or the need for additional monitoring.

The depth of assessment should reflect the examination and the patient's clinical condition. Routine radiography generally requires fewer precautions than contrast-enhanced imaging, procedures involving sedation, or image-guided interventions.

Relevant information may include the reason for examination, current symptoms, medical history, previous imaging experiences, allergies, medication use, renal function where indicated, pregnancy-related considerations, implanted devices, and the patient's ability to tolerate positioning.

Medical Nurses contribute through clinical assessment and identification of patients requiring additional assistance. Radiology professionals evaluate modality-specific risks and determine appropriate protocols.

Pharmacy involvement becomes important when medication-related concerns could affect procedural safety or preparation.

Medical Secretaries help ensure that required documents and approved preparation instructions are available before the appointment.

Health Information systems should support accurate retrieval of relevant clinical information without requiring patients to repeatedly provide details already documented in their records.

An effective assessment process is neither unnecessarily burdensome nor superficial. It should identify clinically meaningful risks while avoiding delays caused by routine testing or restrictions that are not indicated.

6. RADIATION PROTECTION AND OPTIMIZATION OF IMAGING EXPOSURE

Ionizing radiation remains an important consideration in radiography, computed tomography, fluoroscopy, and certain other imaging procedures. Although these examinations provide substantial clinical benefits, exposure should be justified and optimized according to the diagnostic objective.

Radiation protection involves selecting appropriate imaging protocols, avoiding unnecessary repeat examinations, and using technical parameters that achieve adequate diagnostic quality without excessive exposure.

Children require particular attention because of their developing tissues and potentially longer remaining lifespan. Patients undergoing repeated imaging for chronic conditions may also benefit from careful review of cumulative examination history and the clinical necessity of additional studies.

Radiology professionals have primary responsibility for applying radiation protection principles and optimizing technical protocols.

Health Information systems can support access to previous examinations and relevant radiation dose information where available.

Medical Nurses contribute by preparing patients, explaining procedures, and helping reduce movement or positioning difficulties that might otherwise require repeat imaging.

Medical Secretaries may assist with obtaining previous examination records and coordinating appropriate appointments.

Patient communication should present radiation risks in proportion to the expected clinical benefits. Exaggerated warnings may discourage necessary investigations, while inadequate explanations may undermine informed decision-making.

7. CONTRAST MEDIA SAFETY AND INDIVIDUALIZED RISK ASSESSMENT

Contrast agents improve visualization of anatomical structures and physiological processes in many CT, MRI, and other imaging examinations. Their use requires consideration of the specific agent, examination indication, previous reactions, and relevant patient characteristics.

Iodinated contrast media and gadolinium-based contrast agents have different safety profiles and should not be treated as interchangeable.

A previous reaction to a particular contrast agent may be clinically relevant when planning another examination. The nature and severity of the reaction, the agent involved, and available alternatives should be reviewed.

Renal function assessment should be guided by the patient's risk factors, clinical condition, the contrast agent, and applicable evidence-based protocols rather than applied indiscriminately to every patient.

Radiology professionals determine contrast protocols and evaluate imaging-related risk. Pharmacy supports medication review and provides pharmacological expertise when specific concerns arise.

Medical Nurses assess patients, assist with authorized administration where applicable, and monitor for unexpected reactions.

Health Information systems should preserve accurate records of previous contrast exposure and clinically relevant adverse reactions.

Medical Secretaries support appointment coordination when additional assessment or preparation is required, without independently interpreting clinical risk.

The aim is to preserve access to beneficial imaging while identifying and reducing preventable complications.

8. MEDICATION RECONCILIATION AND PHARMACOLOGICAL PREPARATION

Medication reconciliation is particularly important when imaging procedures require fasting, sedation, intravenous contrast, or temporary modification of usual treatment.

Patients may receive medicines for diabetes, hypertension, cardiovascular disease, epilepsy, psychiatric conditions, chronic pain, and other disorders. Some also use over-the-counter products or supplements that are not consistently documented.

A complete medication history helps identify treatment that should be continued, medicines requiring individualized review, and potential interactions with drugs administered during procedures.

Pharmacists can assess the medication regimen and communicate recommendations to the responsible clinical team. Medical Nurses confirm relevant medication information during patient assessment and observe for clinical effects.

Medication adjustments should be based on current evidence and the patient's condition. For example, instructions concerning diabetes medicines during fasting should account for the risk of hypoglycemia and the anticipated duration of reduced oral intake.

Similarly, decisions about anticoagulant or antiplatelet therapy are particularly important for image-guided invasive procedures and must balance bleeding risk against the risk of thrombosis.

Medical Secretaries can communicate approved preparation instructions but should refer medication questions to qualified clinicians.

Accurate Health Information records help prevent conflicting instructions and support continuity when patients attend multiple departments.

9. PREVENTION AND MANAGEMENT OF CONTRAST-RELATED ADVERSE REACTIONS

Although many contrast-enhanced examinations are completed without complications, healthcare teams must remain prepared to recognize and respond to acute adverse reactions.

Reactions may range from mild symptoms to potentially life-threatening events requiring immediate medical treatment.

The clinical team should distinguish contrast-related reactions from other causes of deterioration, including anxiety, vasovagal episodes, or underlying illness.

Medical Nurses and Radiology personnel are often the first to observe symptoms during or shortly after contrast administration. Their responsibilities include recognizing abnormal clinical changes, activating the appropriate emergency response, and providing care within their professional scope.

Pharmacy contributes through ensuring the availability, correct storage, and safe use of emergency medications, as well as supporting review of treatment protocols.

Health Information professionals help ensure that significant reactions are documented accurately and made available for future clinical encounters.

After an event, the record should identify the contrast agent, observed symptoms, timing, treatment, and clinical outcome where known.

Accurate documentation is essential because a vague entry such as “contrast allergy” may not provide sufficient information for future examination planning.

10. MRI SAFETY AND SCREENING FOR IMPLANTED DEVICES

MRI does not use ionizing radiation, but its magnetic and radiofrequency environments create safety considerations that differ substantially from those associated with radiography or CT.

Certain implanted devices, metallic foreign bodies, monitoring equipment, and other objects may be unsafe or subject to specific conditions within the MRI environment.

Appropriate screening is therefore essential before patients, staff, or accompanying individuals enter controlled MRI areas.

Radiology professionals with appropriate MRI safety responsibilities must verify device information and determine whether the planned examination can proceed under the applicable conditions.

Medical Nurses can assist in identifying relevant medical history and ensuring that patients understand screening questions.

Health Information systems should make implant records and previous relevant documentation available whenever possible.

Medical Secretaries may coordinate the collection of device information before appointments, helping reduce avoidable cancellations.

Pharmacy may become involved when MRI preparation includes medication management, sedation, or other pharmacological considerations.

Uncertainty about device compatibility should not be resolved through assumptions or incomplete patient recollection. Verification through reliable device documentation and established MRI safety procedures is necessary.

11. SAFE SEDATION AND ANALGESIA DURING IMAGING PROCEDURES

Some patients cannot comfortably complete an imaging examination without additional support. Severe anxiety, claustrophobia, pain, involuntary movement, developmental limitations, or prolonged procedure duration may create difficulties.

Sedation or analgesia may be appropriate in selected circumstances, but these interventions introduce risks that require careful clinical assessment and monitoring.

Before sedation, the responsible team should evaluate the patient's medical condition, relevant medication history, airway and respiratory considerations, fasting requirements where applicable, and expected recovery needs.

Pharmacy supports medication selection, dose verification, interaction assessment, and safe preparation.

Medical Nurses with appropriate training contribute to monitoring, recognition of complications, and recovery assessment.

Radiology coordinates procedural requirements and ensures that suitable monitoring and emergency resources are available.

Medical Secretaries assist with scheduling and communication of approved preparation and escort requirements.

Health Information systems support documentation of administered medicines, observations, and discharge criteria.

Sedation should be performed only under appropriate clinical authorization, staffing, monitoring, and institutional protocols.

12. NURSING CARE FOR PATIENTS WITH COMPLEX MEDICAL CONDITIONS

Patients with complex medical conditions often require more than routine imaging preparation.

Individuals with cardiovascular instability, respiratory compromise, severe pain, neurological impairment, or reduced consciousness may need continuous monitoring and coordinated transfer arrangements.

Medical Nurses assess whether the patient can safely tolerate transport, positioning, and the anticipated examination.

They may also identify the need for oxygen support, ongoing intravenous therapy, monitoring equipment, or accompanying clinical personnel.

Radiology professionals must understand these requirements when selecting examination protocols and arranging equipment.

Pharmacy contributes when ongoing medicines, infusions, or procedural drugs require review.

Health Information supports the transfer of accurate clinical information, while Medical Secretaries may assist with logistical coordination for planned examinations.

Safe imaging in medically complex patients requires balancing diagnostic urgency against the risks of transport and procedural interruption.

13. PEDIATRIC IMAGING SAFETY AND FAMILY-CENTERED COMMUNICATION

Children present distinctive imaging challenges because of developmental differences, communication needs, and the importance of minimizing unnecessary radiation exposure.

Young patients may be frightened by unfamiliar equipment, separation from caregivers, loud noises, or the need to remain still.

A family-centered approach can improve cooperation by explaining the procedure in age-appropriate language and involving caregivers when appropriate.

Radiology professionals select pediatric imaging protocols and determine whether alternatives can provide the required diagnostic information.

Medical Nurses support preparation, comfort, and monitoring when needed.

Pharmacy contributes to weight-appropriate medication assessment and the safe use of procedural medicines where indicated.

Medical Secretaries help families understand appointment arrangements and approved preparation instructions.

Health Information systems must maintain accurate age, weight, medical history, and previous imaging records.

The objective is to obtain diagnostically useful images while reducing avoidable distress, unnecessary exposure, and repeated examinations.

14. DIAGNOSTIC IMAGING IN OLDER ADULTS AND PATIENTS WITH FRAILTY

Older adults frequently present with combinations of chronic disease, reduced mobility, sensory impairment, cognitive difficulties, and polypharmacy.

These factors can affect examination preparation, communication, transport, positioning, and recovery.

A patient with frailty may experience considerable discomfort when required to remain in a fixed position. Someone with cognitive impairment may struggle to understand instructions or tolerate an unfamiliar environment.

Medical Nurses can identify functional limitations and coordinate assistance. Radiology professionals may adapt positioning and examination techniques while preserving diagnostic quality.

Pharmacy can review medications that may contribute to sedation, confusion, falls, or other clinically relevant risks.

Medical Secretaries support appointment arrangements that account for transport and caregiver availability.

Health Information systems help ensure that previous clinical assessments and relevant medical details are accessible.

A patient-centered approach avoids treating chronological age alone as a contraindication to imaging. Decisions should reflect the individual's clinical condition, functional status, and expected benefit.

15. IMAGING SAFETY DURING PREGNANCY AND BREASTFEEDING

Diagnostic imaging during pregnancy requires careful consideration of clinical necessity, examination type, and the potential consequences of delaying diagnosis.

Ultrasonography and MRI without contrast may be useful alternatives in some circumstances, but examinations involving ionizing radiation may still be appropriate when clinically justified.

The relevant risks depend on the imaging modality, body region, examination parameters, and stage of pregnancy.

Radiology professionals are responsible for selecting appropriate protocols and discussing imaging-related considerations with the referring team.

Medical Nurses assist with clinical assessment and communication.

Pharmacy may provide guidance when contrast agents, sedation, or other medicines are involved.

Health Information systems should maintain accurate documentation of relevant clinical details and informed discussions.

Medical Secretaries can facilitate timely appointments and direct pregnancy-related questions to qualified clinical staff.

Patients should receive balanced information that supports necessary diagnosis without creating avoidable fear.

16. INFECTION PREVENTION AND SAFE PATIENT MOVEMENT

Diagnostic imaging departments receive patients from emergency units, inpatient wards, intensive care facilities, and outpatient services. This movement creates opportunities for transmission of infection when appropriate precautions are not maintained.

Infection prevention requires effective cleaning of equipment and examination surfaces, hand hygiene, appropriate personal protective equipment, and procedures suited to the patient's infection risk.

Medical Nurses contribute by communicating relevant precautions and supporting safe transport and patient handling.

Radiology personnel implement examination-area infection control measures and coordinate equipment preparation.

Medical Secretaries may assist with scheduling arrangements that reduce unnecessary crowding and support appropriate patient flow.

Health Information systems enable authorized staff to access relevant infection-control information.

Pharmacy supports the safe management of medicines and products used during imaging-related procedures.

Infection prevention should be integrated into routine workflow rather than treated as a separate activity performed only after an exposure or outbreak.

17. COMMUNICATION OF CRITICAL AND UNEXPECTED IMAGING FINDINGS

A diagnostic examination may identify findings requiring urgent clinical intervention, even when the original referral concerned a different suspected condition.

Examples include intracranial hemorrhage, major vascular abnormalities, or other findings requiring immediate assessment.

Timely reporting alone may be insufficient if the responsible clinical team does not receive or acknowledge the information.

Radiology departments should maintain clear procedures for communicating critical results directly to appropriate clinicians.

Health Information systems can support electronic alerts, documentation of communication, and tracking of acknowledgment where available.

Medical Nurses may facilitate urgent clinical reassessment when informed of significant findings.

Medical Secretaries may assist with locating appropriate contacts or coordinating follow-up under approved procedures, but should not independently interpret results or determine clinical urgency.

A reliable process should include escalation when the intended recipient cannot be reached.

Closed-loop communication is especially important because the clinical consequences of a delayed response may be substantial.

18. REDUCING MISSED APPOINTMENTS AND INCOMPLETE DIAGNOSTIC FOLLOW-UP

Missed imaging appointments and incomplete follow-up can delay diagnosis, prolong uncertainty, and lead to repeated clinical visits.

The reasons may include unclear instructions, financial or transport difficulties, language barriers, scheduling conflicts, or limited understanding of the examination's importance.

Medical Secretaries are central to appointment coordination and communication with patients.

Health Information systems can support tracking of pending examinations, cancellations, completed studies, and outstanding reports.

Medical Nurses may identify patients requiring additional preparation or clinical assistance, while Pharmacy can help resolve medication-related barriers.

Radiology should provide clear examination requirements and clinical prioritization.

When an examination is missed, follow-up procedures should distinguish routine scheduling issues from potentially urgent clinical concerns.

Patients with abnormal findings requiring further investigation may need additional coordination to prevent loss to follow-up.

19. DIGITAL INTEGRATION OF RADIOLOGY AND CLINICAL INFORMATION SYSTEMS

Digital information systems have transformed the organization of diagnostic imaging services.

Radiology information systems, picture archiving and communication systems, electronic health records, and electronic referral platforms can improve access to examination details, clinical histories, images, and reports.

However, digital integration also introduces risks when systems contain incomplete information, inconsistent patient identifiers, delayed updates, or poorly designed communication workflows.

Health Information professionals contribute through data governance, record integrity, access management, and information exchange.

Radiology professionals rely on accurate digital information for protocol selection, comparison with previous studies, and reporting.

Medical Nurses and Pharmacy professionals require access to relevant clinical information when preparing and monitoring patients.

Medical Secretaries use administrative systems to coordinate appointments and maintain referral records.

Effective interoperability should reduce unnecessary duplication while ensuring that sensitive information remains accessible only to authorized personnel.

20. PATIENT EDUCATION, INFORMED PARTICIPATION, AND HEALTH LITERACY

Patients are more likely to complete imaging examinations successfully when they understand what will happen and why particular preparation is necessary.

Unclear instructions can lead to anxiety, missed appointments, inappropriate fasting, medication errors, or failure to attend recommended follow-up.

Education should be tailored to the examination and the patient's ability to understand the information.

Medical Nurses can explain clinical preparation and recovery requirements. Radiology personnel describe procedural expectations and modality-specific considerations.

Pharmacy provides individualized medication counseling when needed.

Medical Secretaries communicate approved appointment instructions and help patients obtain clarification from clinical staff.

Health Information systems may support written instructions, patient portals, and access to examination reports.

Patients should have opportunities to ask questions and communicate concerns before the examination.

The quality of patient education should be assessed by whether patients understand and can act on the information, not merely whether written instructions were provided.

21. PRIVACY, CONFIDENTIALITY, AND RESPECT FOR PATIENT DIGNITY

Diagnostic imaging frequently involves sensitive clinical information and, in some examinations, exposure of private areas of the body or discussion of potentially serious diagnoses.

Maintaining patient dignity requires appropriate physical privacy, respectful communication, and careful handling of clinical information.

Health Information professionals support confidentiality through secure records, appropriate access permissions, and information governance.

Radiology and Nursing staff should ensure that examinations are performed with suitable privacy measures and that patients receive clear explanations before positioning or procedures.

Medical Secretaries must protect personal information during registration, scheduling, and communication.

Pharmacy professionals have similar responsibilities when discussing medicines and clinical conditions.

Patient-centered privacy extends beyond preventing unauthorized disclosure. It also involves avoiding unnecessary exposure, respecting cultural and personal preferences, and maintaining professional sensitivity throughout the imaging encounter.

22. MANAGING DIAGNOSTIC ERRORS AND PREVENTABLE IMAGING INCIDENTS

Diagnostic errors may arise from inappropriate examination selection, technical limitations, incomplete clinical information, interpretation problems, or failure to communicate and act upon results.

Not every diagnostic discrepancy represents preventable error, and uncertainty is an inherent part of clinical medicine.

Nevertheless, healthcare institutions should investigate incidents where system weaknesses may have contributed to avoidable harm.

Radiology can review protocol selection, image quality, reporting practices, and communication procedures.

Health Information professionals support accurate incident records and analysis of documentation or information-transfer failures.

Medical Nurses and Pharmacy professionals provide insight into clinical and medication-related circumstances.

Medical Secretaries may identify referral, scheduling, or communication problems that affected the examination pathway.

A constructive review process should distinguish individual performance concerns from broader system failures and focus on improvements that reduce recurrence.

23. INTERPROFESSIONAL EDUCATION AND COMPETENCY DEVELOPMENT

Safe diagnostic imaging depends on professionals understanding both their own responsibilities and the contributions of other departments.

Training should address patient identification, clinical preparation, contrast safety, MRI screening, radiation protection, medication management, communication of critical results, and emergency response.

Interprofessional education can help clarify where responsibilities overlap and where clinical decisions require escalation.

For example, a simulation involving a patient with a previous severe contrast reaction may require coordinated action from Radiology, Nursing, Pharmacy, administrative personnel, and Health Information staff.

Such exercises can reveal gaps in communication or documentation that may not be apparent during routine operations.

Training should be adapted to professional scope and institutional responsibilities.

Medical Secretaries do not require the same clinical competencies as Nurses or Pharmacists, but they do need to recognize when a question or incomplete document must be referred to qualified personnel.

Regular education supports consistency as imaging technologies, clinical guidance, and organizational systems evolve.

24. QUALITY INDICATORS AND CONTINUOUS IMPROVEMENT IN DIAGNOSTIC IMAGING

Quality improvement requires meaningful measurement of both technical performance and the wider patient experience.

Relevant indicators may include wrong-patient or wrong-examination events, preventable repeat imaging, contrast-related incidents, incomplete screening, appointment cancellations due to inadequate preparation, reporting delays, and failures in communication of critical findings.

Patient-centered measures may include waiting times, accessibility, understanding of preparation instructions, comfort during examinations, and satisfaction with communication.

Health Information professionals can support data collection, validation, and reporting.

Radiology, Pharmacy, Nursing, and administrative teams should participate in interpreting findings and identifying practical improvements.

Performance measures must be used carefully. A higher number of reported safety incidents may reflect improved reporting rather than worsening practice.

Quality improvement should therefore examine incident severity, reporting completeness, process reliability, and whether corrective actions produce sustained benefits.

25. PROPOSED COLLABORATIVE MODEL FOR COMPREHENSIVE IMAGING SAFETY

A practical multidisciplinary model can be organized around six connected stages: referral review, preparation, risk assessment, examination, result communication, and follow-up.

During referral review, Radiology evaluates examination appropriateness and protocol requirements, while Medical Secretaries ensure that necessary administrative information is complete.

During preparation, Medical Nurses assess clinical needs, Pharmacy addresses relevant medication concerns, and patients receive clear instructions.

Risk assessment incorporates modality-specific screening, review of available clinical information, and identification of patients requiring additional precautions.

During the examination, Radiology personnel maintain technical and procedural safety, with Nursing and other clinical support according to patient needs.

After image acquisition, reporting and communication systems should ensure that clinically important findings reach the responsible care team.

Finally, Health Information systems and administrative coordination support documentation, follow-up scheduling, and continuity of care.

The effectiveness of this model depends on shared accountability without confusion about professional authority.

Each department contributes a different form of expertise, but responsibility for clinical decisions must remain with appropriately qualified professionals.

26. FUTURE DIRECTIONS IN PATIENT-CENTERED DIAGNOSTIC IMAGING

The future of diagnostic imaging will be shaped by increasingly sophisticated technology, expanding use of artificial intelligence, greater integration of electronic health records, and rising expectations for personalized healthcare.

These developments may improve image acquisition, workflow prioritization, clinical decision support, and identification of patients at risk of delayed follow-up.

However, technological progress will not eliminate the need for careful clinical judgment, clear communication, and coordinated patient care.

Artificial intelligence applications in imaging require evaluation for accuracy, bias, transparency, and their effects on clinical workflow. Automated alerts and decision-support tools should assist qualified professionals rather than replace their responsibilities.

Health Information teams will play an increasingly important role in data quality, system interoperability, privacy, and governance.

Radiology professionals will remain responsible for appropriate clinical interpretation and procedural oversight, while Pharmacy and Nursing will continue to address medication safety and patient-specific clinical needs.

Medical Secretaries will remain important in helping patients navigate increasingly complex diagnostic services, particularly when examinations involve multiple appointments, specialized preparation, or follow-up requirements.

Future research should examine how coordinated departmental practices influence preventable imaging incidents, diagnostic delays, patient understanding, and the completion of recommended care.

The broader objective is to establish imaging services in which technological accuracy is supported by reliable clinical preparation, effective information exchange, and sustained attention to the patient's experience.

CONCLUSION

Comprehensive diagnostic imaging safety extends beyond the technical performance of radiological examinations. It requires careful patient assessment, appropriate examination selection, safe medication practices, effective communication, accurate documentation, and timely follow-up. As imaging services become increasingly complex, fragmented responsibilities can contribute to preventable complications, diagnostic delays, and gaps in continuity of care.

Collaboration among Radiology, Pharmacy, Medical Nurse, Medical Secretary, and Health Information professionals provides an important foundation for addressing these challenges. Radiology ensures diagnostic appropriateness and procedural safety, Pharmacy contributes medication-related expertise, Medical Nurses support patient preparation and clinical monitoring, Medical Secretaries coordinate appointments and essential communication, and Health Information professionals maintain reliable clinical records and information exchange. When these responsibilities are clearly defined and coordinated, patients are better supported throughout the diagnostic pathway.

Patient-centered imaging also requires attention to individual circumstances, including age, comorbidities, functional limitations, health literacy, psychological concerns, and personal preferences. Standardized safety procedures, effective communication of critical findings, appropriate use of digital systems, and continuous quality improvement can strengthen diagnostic reliability while reducing unnecessary delays and avoidable harm.

Ultimately, the quality of diagnostic imaging should be measured not only by the accuracy of the images produced but also by how safely, efficiently, and compassionately patients move from referral to diagnosis and subsequent care. Sustained collaboration across these five departments can help healthcare institutions deliver imaging services that are clinically effective, operationally reliable, and responsive to patients' needs.

REFERENCES

[1] World Health Organization. *Global patient safety action plan 2021–2030: towards eliminating avoidable harm in health care*. Geneva: World Health Organization; 2021.

[2] World Health Organization. *Radiation protection and safety in medical uses of ionizing radiation*. Specific Safety Guide No. SSG-46. Vienna: International Atomic Energy Agency; 2018.

[3] International Commission on Radiological Protection. The 2007 recommendations of the International Commission on Radiological Protection. ICRP Publication 103. *Ann ICRP*. 2007;37(2–4):1-332.

[4] International Commission on Radiological Protection. Radiological protection in medicine. ICRP Publication 105. *Ann ICRP*. 2007;37(6):1-63.

[5] International Commission on Radiological Protection. Diagnostic reference levels in medical imaging. ICRP Publication 135. *Ann ICRP*. 2017;46(1):1-144.

[6] American College of Radiology. *ACR manual on contrast media*. Reston (VA): American College of Radiology; 2024.

[7] European Society of Urogenital Radiology. *ESUR guidelines on contrast agents*. Version 10.0. Vienna: European Society of Urogenital Radiology; 2018.

[8] Davenport MS, Perazella MA, Yee J, Dillman JR, Fine D, McDonald RJ, et al. Use of intravenous iodinated contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology*. 2020;294(3):660-668.

[9] Weinreb JC, Rodby RA, Yee J, Wang CL, Fine D, McDonald RJ, et al. Use of intravenous gadolinium-based contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology*. 2021;298(1):28-35.

[10] American College of Radiology. *ACR manual on MR safety*. Reston (VA): American College of Radiology; 2024.

[11] Kanal E, Barkovich AJ, Bell C, Borgstede JP, Bradley WG Jr, Froelich JW, et al. ACR guidance document on MR safe practices: 2013. *J Magn Reson Imaging*. 2013;37(3):501-530.

[12] American College of Radiology. *ACR appropriateness criteria*. Reston (VA): American College of Radiology.

[13] European Society of Radiology. ESR paper on structured reporting in radiology. *Insights Imaging*. 2018;9(1):1-7.

[14] Berlin L. Communicating findings of radiologic examinations: whither goest the radiologist's duty? *AJR Am J Roentgenol*. 2002;178(4):809-815.

[15] Brady AP. Error and discrepancy in radiology: inevitable or avoidable? *Insights Imaging*. 2017;8(1):171-182.

[16] Brady AP, Laoide RÓ, McCarthy P, McDermott R. Discrepancy and error in radiology: concepts, causes and consequences. *Ulster Med J*. 2012;81(1):3-9.

[17] European Society of Radiology. Patient safety in medical imaging: a joint paper of the European Society of Radiology and the European Federation of Radiographer Societies. *Insights Imaging*. 2019;10:45.

[18] Institute of Medicine. *Improving diagnosis in health care*. Washington (DC): National Academies Press; 2015.

[19] Kripalani S, LeFevre F, Phillips CO, Williams MV, Basaviah P, Baker DW. Deficits in communication and information transfer between hospital-based and primary care physicians: implications for patient safety and continuity of care. *JAMA*. 2007;297(8):831-841.

[20] Mueller SK, Sponsler KC, Kripalani S, Schnipper JL. Hospital-based medication reconciliation practices: a systematic review. *Arch Intern Med*. 2012;172(14):1057-1069.

[21] Mekonnen AB, McLachlan AJ, Brien JAE. Pharmacy-led medication reconciliation programmes at hospital transitions: a systematic review and meta-analysis. *J Clin Pharm Ther*. 2016;41(2):128-144.

[22] Agency for Healthcare Research and Quality. *Diagnostic safety and quality*. Rockville (MD): Agency for Healthcare Research and Quality.

[23] World Health Organization. *Medication without harm: WHO global patient safety challenge*. Geneva: World Health Organization; 2017.

[24] World Health Organization. *Patient safety: making health care safer*. Geneva: World Health Organization; 2017.

[25] European Society of Radiology. ESR statement on the holistic approach to patient care in radiology. *Insights Imaging*. 2019;10:18.