

Original Research

View Article Online



Received 11 December 2022

Revised 14 April 2023

Accepted 18 August 2023

Available online 29 August 2023

Edited by Si Mi

KEYWORDS:

Residue variability
Pesticide toxicity
Agricultural workers
Pesticide safety
Risk assessment
Exposure assessment
Pesticide residues
Residue dissipation
Cereal grains

Natr Resour Human Health 2023; 3 (4): 387-402
<https://doi.org/10.53365/nrfhh/171368>
eISSN: 2583-1194
Copyright © 2023 Visagaa Publishing House

Concentrations of Pesticides Residues in Grain Sold at Selected Markets of Southwest Nigeria

Modupe Abeke Oshatunberu¹, Adebayo Oladimeji², Sawyerr Olawale Henry¹, Opasola Afolabi Olaniyan¹, Morufu Olalekan Raimi^{1, 3,*}

¹Department of Environmental Health, Faculty of Pure and Applied Science, Kwara State University, Malete, Kwara State. Nigeria

²Department of Zoology, Faculty of Pure and Applied Science, Kwara State University, Malete, Kwara State. Nigeria

³Department of Environmental Management and Toxicology, Faculty of Science, Federal University Otuoke, Bayelsa State, Nigeria

ABSTRACT: The objective of this work is to determine the concentration of the identified pesticide in grains commonly used by farmers, and which are available directly from the open markets in the Nigerian market. Pesticide residues were quantified through a multiresidue method using a varian 3800/4000 Gas Chromatograph Mass Spectrometer (GC/MS). The simultaneous measurement of four classes of pesticides by gas chromatography-mass spectrometry (GC/MS) approach using sample preparation on QuEChERS-citrate, was developed and validated. The results frequently showed high specific contamination, which makes sense given the target market and dietary diversity in the area. This study found that grains purchased from particular markets in southwest Nigeria contained numerous pesticide residues. The MRLs set by the EU or FAO/WHO or both were surpassed by 17 out of the total 27 pesticides reported in this work in at least one grain, despite the fact that there were no published codex MRLs for some pesticide residues in some grains. These residues were classified among the four pesticides classes: carbamates, organophosphates, organochlorines, and pyrethroids. In actuality, 90% of the mainly banned organochlorine pesticides exceeded MRLs. Thus, this study revealed higher concentration levels of organo-chloride pesticides and organo-phosphate pesticides in grain samples drawn from selected markets in southwest Nigeria. Consumers seeking high-quality food in Nigeria should take note of these facts. Notwithstanding, the small number of samples with detectable residues suggests that there is a necessity to increase monitoring of pesticides in grains, educating farmers, and raising their awareness of the dangers associated with the unauthorized use of pesticides that are only allowed in agriculture and can damage the reputation of the sector as a whole.

1. INTRODUCTION

Regardless of where you live or who you are, food safety is crucial for human health. Consumers in Nigeria are commonly exposed to pesticide residues through their food. What consequences, if any, such chemical pollutants cause to consumer health remain unclear given the presence of pesticide residues in food. While, people who live in rural areas are frequently exposed to pesticides, solvents, and toxic elements, as well as other environmental risks linked to agricultural output. According to reports, chronic exposure to pesticides can cause conditions like diabetes, cancer, cardiovascular disease, and neurological issues (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr,

et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; M.R. Olalekan et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Consequently, there is now much greater awareness about the health concerns associated with exposure to pesticides (Asiegbu et al., 2022; O.R. Morufu, 2021,?; O.R. Morufu, Tonye, et al., 2021; M.R. Olalekan et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). With the global population anticipated to reach about 10 billion people by 2050, increasing food production is every nation's top priority (M.O. Raimi, Bilewu, et al., 2019; M.O. Raimi, Suleiman, et al., 2019). Evidence suggests that the global population is growing by 97 million people yearly. In fact, the Food and Agricultural Organization (FAO) of the United Nations has made the

* Corresponding author.

E-mail address: morufuolalekan.raimi@gmail.com (Morufu Olalekan Raimi)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

grim prediction that in order to meet the demand of a growing population, global food production must expand by 70% (R.M. Olalekan et al., 2019; M.O. Raimi, Abiola, et al., 2022). However, increasing food production is beset by a number of increasingly difficult problems, not the least of which is the extremely limited expansion of cultivable land. Because of this, the existing agricultural system is under immense pressure to produce enough food using the same amount of land, water, and other resources as before (Suleiman et al., 2019). Insecticides, herbicides, nematicides, fungicides, fertilizers, and soil amendments are currently utilized in greater amounts than in the past in order to increase crop production (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Since the development of synthetic insecticides in 1940, when organochlorine (OCl) pesticides were first applied for pest control, these chemicals have mostly entered the scene. Prior to this introduction, the majority of weeds, pests, insects, and illnesses were managed utilizing environmentally friendly techniques such cultural, mechanical, and physical control methods (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Pesticides are now a crucial part of modern life and are used to preserve food storage facilities, agricultural land, flower gardens, as well as to get rid of pests that spread infectious diseases that are dangerous to humans (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020; M.O. Raimi & Raimi, 2020; Samson et al., 2020). Pesticides are thought to cost roughly \$38 billion worldwide each year (Pan-Germany, 2012). To address the need on a global scale, manufacturers and researchers are developing novel pesticide formulations. Ideally, the pesticides used should only be poisonous to the organisms they are intended to kill, be somewhat biodegradable, and environmentally beneficial (Rosell et al., 2008; Suleiman et al., 2019). Sadly, this is seldom the case because the majority of non-selective pesticides are general-purpose and may harm organisms that are beneficial to the ecosystem or harmless. The majority of pesticides are thought to poison the environment, with just around 0.1% of them reaching the intended target organisms (Asiegbu et al., 2022; Carriger et al., 2006; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). The ongoing use of pesticides that are persistent and non-biodegradable has damaged a variety of ecosystem components in the water, air, and soil (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). In the upper tropic level of the food chain,

pesticides have also bioaccumulated (M.O. Raimi, Olawale, et al., 2022; Suleiman et al., 2019). More recent times, a number of acute and chronic health illnesses have been connected to pesticide exposure (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; Mostafalou & Abdollahi, 2012; M.R. Olalekan et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Many people are well aware that pesticide chemicals frequently come into contact with food residues (R.M. Olalekan et al., 2020). Since pesticides are frequently utilized due to their capacity to destroy biological species like insects, plant diseases, and weeds, many consumers may find this knowledge to be upsetting. The topic of what risks, if any, consumers are exposed to as a result of consuming pervasive pesticide residues in their diet is raised by their existence in food. Advocates frequently give advice to consumers regarding safety issues, such as whether to eat conventional or organic foods, which geographic sources of foods should be avoided, and what may be done to decrease exposure to pesticides by washing, peeling, boiling, or scrubbing foods before consumption (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Such recommendations usually lack the necessary scientific support. Since pesticides, like all chemicals, adhere to the rules of toxicology, the mere presence of a pesticide residue on a food item does not prove that it has caused harm. The potential dangers of a chemical are determined by its dose or exposure from series of studies (Afolabi & Morufu, 2021; Afolabi & Raimi, 2021; Mo & Sabinus, 2017; Odipe et al., 2018; Omotoso et al., 2021; M.O. Raimi et al., 2021; O. Raimi et al., 2022; Stephen et al., 2022). Contrary to the wise counsel from healthcare professionals that calls for balanced and diverse diet rich in whole grains, consumers may be opting to heed dubious advice to stay away from particular foods because of possible pesticide residues. Studies show that residues are frequently found, and that people regularly come into contact with pesticides in their food (Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020). It is frequently implied that pesticide residue discoveries that contravene National Agency for Food and Drug Administration and Control (NAFDAC) food safety standards may endanger the health of consumers because the relationship between permissible levels of pesticides on foods (tolerances) and safety is poorly understood (R.M. Olalekan et al., 2020). Despite having a population of 213 million, Nigeria ranks fourth in the world for the number of undernourished individuals Ogisi and Begho (2021). Due to their high nutritious content, staple foods including maize, rice, millet, and cowpea present a chance to reduce malnutrition. For instance, compared to animal products, cowpeas are a cheap source of protein whereas maize is high in carbohydrates (James et al., 2020). However, malnutrition has been associated with relying too heavily on one or a few of these basics (Ekpa et al., 2018). In order to optimize mineral bioavailability and fortification,

it would be helpful to understand the nutritional makeup of these grains. The environment, animal products, cash crops, vegetables, and processed foods have been the main subjects of the most recent studies on pesticide residues in food in Nigeria. Additionally, research on cereals like maize has only been done in a confined region (Sosan et al., 2018). A thorough analysis of pesticide residues in common foods in various states is required to determine potential dangers and exposure and to protect the general public's health in light of the requirement for regular monitoring of toxins in food (Adiama et al. (2022); Habeeb et al. (2022); H.M. Lateefat et al. (2022); M.H. Lateefat, Opasola, Adiama, et al. (2022); M.H. Lateefat, Opasola, Misbahu, and Morufu (2022)). Even though the processes for the approval of pesticide active substances for agricultural use adhere to strict regulations to ensure both their high effectiveness and their environmental and consumer safety (Afolabi & Morufu, 2021; Afolabi & Raimi, 2021; Mo & Sabinus, 2017; O.R. Morufu, Clinton, & Bowale, 2021; O.R. Morufu, Henry, et al., 2021; O.R. Morufu, Olawale, & Clinton, 2021; R. Morufu & E, 2017; Odipe et al., 2018; A.S. Olalekan et al., 2022; M.R. Olalekan, Albert, et al., 2022; M.R. Olalekan, Olawale, et al., 2022; R.M. Olalekan et al., 2018; Omotoso et al., 2021; M. Raimi & Sawyerr, 2022; M.O. Raimi et al., 2021; M.O. Raimi, Sawyerr, et al., 2022; O. Raimi et al., 2022; Stephen et al., 2022), there are significant concerns about the health risks associated with exposure to many pesticides while working as well as consumer exposure to their residues in food and drinking water (R.M. Olalekan et al., 2020). The accumulation of pesticide residues in agricultural crops is commonly linked to the irrational use of pesticides, which can have negative effects on both the environment and human health in the short and long term (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Numerous pesticides and fungicides used to control crop diseases and pests are now known to be carcinogenic, harmful to developing fetuses, endocrine disruptors, and to increase the risk of immunological, neurological, and respiratory illnesses (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Numerous research has confirmed the link involving pesticide usage and an elevated likelihood of different malignancies as well as other health issues (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Young infants are particularly at risk from pesticide exposure connected to various sources, making them susceptible to possible neurological, neurodevelopmental, as well as other consequences (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). The

development of a child's brain nor central nervous system can be hampered by even minor early-life pesticide exposure (Asiegbu et al., 2022). Consumers who desire food free of agricultural residues are concerned about the hazards from exposure to pesticides that are all mentioned above and others (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; M.R. Olalekan et al., 2021; R.M. Olalekan et al., 2020). The majority of consumers believe that grains are safe and good for their health, and they link this belief to the lack of agricultural chemical residues, as well as to a natural flavor and concern for the environment. Its restricted production practices are the cause of these perceived qualities, which increase consumer trust across the globe. In addition, a significant and essentially constant proportion of food products with detectable pesticide residues are present on the European market, according to reports on the level of pesticide contamination in food. The tests conducted yearly as part of the mandated control programs encompassing the entire EU and affiliated nations have brought this issue to light in recent years. The European Food Safety Authority (EFSA) is in charge of this market surveillance throughout Europe. According to the EFSA annual reports, between 39.5 and 45.5% of the tested samples had detectable pesticide residues in recent years. Additionally, between 2.6 and 5.1% of the samples included residues that were above the permitted limits (referred to as the MRL-maximum residue level in food of plant origin). Nevertheless, EFSA monitoring shows that a portion of organic samples contain detectable pesticide residues even though the use of synthetic pesticides in organic agriculture is explicitly prohibited by EU Regulation 2018/848. 8.3% of organic samples had detectable pesticide residues in 2015, and 19.9% did so in 2020. Of the analyzed organic samples, 0.7% had residues above the MRL in 2015, and 1.5% did so in 2020. When purchasing grains, consumers have a right to anticipate what they receive will be of good nutritional and biological value and won't be harmful to their health. Farmers must adhere to national and international laws and regulations in order to safeguard consumers and themselves from a variety of risks. Food safety precautions begin with farm production and continue all the way through the supply chain (Hussain et al., 2021a, 2021b; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020). Thus, this paper's objective is to provide and calculate the discovered pesticide's concentration in grains using maximum residue limits (MRL). Consumers, who are increasingly looking for high-quality, pesticide-free grains, may be quite interested in the findings on the safety status of these popular grains originating from the chosen market in the southwest region of Nigeria.

2. MATERIAL AND METHODS

2.1. Study Area

The study area is selected markets across four states (Ekiti, Oyo, Osun and Ondo States) in Southwestern, Nigeria. The markets were: Oja-titun (market) Ile-Ife (16° 18' N 23° 33' E),

Osun; Alesinloye Market Ibadan, Oyo (7° 26' N 3° 55' E; Oja Oba, Ado-Ekiti, Ekiti (7° 37' N 5° 13' E); and Oja Oba, Akure Ondo State (7° 15' N 5° 12' E) (Figure 1). Hereinafter, the market will be referred to as Ekiti, Oyo, Osun and Ondo, for clarity.




Figure 1. Map showing the study area

2.2. Sample collection and preparation

Samples of dry beans, maize, millet, and rice (local and foreign) were purchased from the four markets. Six samples of: brown and white beans, white and yellow maize, brown millet and rice were collected from all four markets except for Ondo (Oja-Oba market, Akure) where yellow maize was not available. The grain samples were bought in the dried state such that there was no need for the samples to undergo additional drying. Patronized grain merchant and purchased each of the six grains sample. Each grain sample was bought in three separate portions of 200 – 250 g and put in black-coloured polyethylene plastic bags, labelled and transported to the laboratory. On arrival, the samples were sorted to remove impurities including stones and shafts. Thereafter, the samples were thoroughly ground using mortar and pestle, and thereafter a hand-grinding machine was used to pulverize into fine powder. Finally, each of the powdered grain sample was stored in labelled Ziploc bag, and kept at 4°C in a refrigerator.

2.3. Proximate analysis

The grain samples were subjected to proximate analysis including ash content, moisture content, crude fat, crude fibre, crude protein, and total carbohydrate (AOAC, 2000). Triplicate samples were constituted from the four markets. The replicates of each market were pooled. The detailed experimental procedure was given as supplementary information (Appendix A).

2.4. Determination of pesticide residues

2.4.1 Chemicals and reagents

Commercial standards of Quinalphos, Diazinon, Fenitrothion, Dimethoate, Acephate, Malathion and Chlorpyrifos were gotten from Sigma Aldrich Laborchemikalien (St Louis, MO, USA) through Top Scientific Ltd. Ado-Ekiti Ekiti state, Nigeria. They all had >99.6% purity. Other reagents including sodium chloride (NaCl), acetone, methanol, anhydrous magnesium sulphate (MgSO₄), HPLC gradient grade acetonitrile and Primary Secondary Amine (PSA) were purchased from Top Scientific Ltd. Ado-Ekiti Ekiti state, Nigeria.

2.4.2 Preparation of pesticide standard solution

The stock solutions of each of the aforementioned commercial standard reagents were prepared separately in acetone (1000 mg/L). A mixed standard solution (50 mg/L acetone) containing the standard reagents was prepared from the individual stock solution and then a lower concentration of 10 mg/L. Thereafter, 0.1, 0.2, 0.5, 1.0, 2.0, 3.0, and 5.0 mg/L working standard were prepared by measuring the required volume to make up each of the concentration in a 10 ml flask and making it up with the stock solution of acetone.

2.4.3 Extraction and clean up

QuEChERS extraction was carried out as described by the European Committee for Standardization (ILNAS-EN, 2018). First, 10 g of the homogenized sample was weighted into a 50 mL polypropylene centrifuge tube, 10 mL of acetonitrile (MeCN) was added followed by the use of a vortex mixer to shake the centrifuge for 30 s. NaCl (1 g) and Anhydrous MgSO₄ (4 g) were added followed by immediate shaking with the vortex mixer for 60 s to pre-empt the MgSO₄ from becoming aggregates and then, the extract was centrifuged at 5000 rpm for 5 minutes. A 3 mL aliquot of the MeCN layer was moved into a 15 mL micro centrifuge tube holding 600 mg and 120 mg of anhydrous MgSO₄ and Primary Secondary Amine (PSA), respectively. The vortex was again used thoroughly for 30 s before centrifuged at 4000 rpm for 5 minutes. Finally, a 0.2 µm PTFE filter was applied in filtering 1 mL supernatant before taking in a clean vial for injection.

2.4.4 Instrumentation and methodology of GC-MS analysis

A Varian 3800/4000 gas chromatograph mass spectrometer was used to analyze pesticide residues. The capillary column was set at AT-1 and a length of 30 m, while the ID and film thickness were set at 0.25 mm 0.25 µm, respectively. The GC initial temperature of the GC column was 70 °C and increase to 300 °C after two minutes holding time. At 300 °C, it was subjected to another 7 minutes holding time. Together, the total run time was 32 minutes. Nitrogen (99.9995% purity) was the carrier gas of choice. The GC-MS interface temperature was at 280 °C and carrier gas had a constant flow rate of 1.51 ml/min. Injector and detector temperatures were set at 250 °C, linear velocity was the flow control mode; and a 1 µl of

sample was injected in the split ratio of 30:0. The range of the MS was 30 – 800 Da. The concentration of pesticide residues in each sample were obtained following the comparative peak retention times of the samples relative to the pure analytical standards; and were represented as relative area percentage. For the identification, the spectrum obtained through GC – MS compounds were compared with the database of the National Institute Standard and Technology. As a precaution, samples were randomly and continuously injected as a batch in order to separate technical variations from the biological medium. More so, the prepared pooled samples were injected into the machine at regular intervals as quality control. The concentrations of the pesticide residues are given in mg/kg.

2.5. Survey

Structured questionnaires were administered to 60 respondents in each of the four markets (Odogwu et al., 2021). The questionnaire included questions relating to the: socio-demographic information, attitudes of grain merchants towards grains preservation, knowledge of grain merchants towards pesticide application and the health implication of pesticide usage on humans Appendix.

2.6. Statistical analyses and calculations

Data analyses were carried out on SPSS version 21.0, IBM, USA. The differences between the treatments were evaluated using Analysis of variance (ANOVA) while Duncan's new multiple-range test was used to separate means at $P < 0.05$. All graphs were plotted on Microsoft Excel, 2019.

3. RESULTS

3.1. Concentration of Identified Pesticide in Beans

3.1.1 Organophosphates (OPPs)

The OPP residues in the bean samples were in the range of 0.014 – 0.036 mg kg⁻¹, 0.014 – 0.042 mg kg⁻¹, 0.010 – 0.014 mg kg⁻¹, 0.011 – 0.045 mg kg⁻¹ and 0.022 – 1.230 mg kg⁻¹ for Phenthoate, Chlorthiophos, Ethion, Prothiofos and Iodofenphos, respectively (Table 1 & 2). Compared to the MRL set by the EU, the level of Ethion in the bean samples was lesser or equal to the stipulated value of 0.01. Only brown beans from Ondo and White beans from Ado-Ekiti were equal to this value.

3.1.2 Organochlorine (OCP)

shows the mean concentrations of OCP residues in beans samples. δ -Lindane and β -Hexachlorocyclohexane were higher than the MRL of 0.01 mg kg⁻¹ set by the FAO/WHO and the EU in both brown and white beans in all the markets. Likewise, the level of p,p'-DDT, Dieldrin were also higher than the MRL in brown beans across all the markets and white beans in Ado-Ekiti, Ibadan and Ondo. While the concentrations of dieldrin in brown beans of all the markets was higher than the MRL set by both the FAO/WHO and the EU, its concentration in brown beans was higher than the FAO/WHO MRL and the EU standard. On the bright side, the concentrations of Aldrin

in the brown and white beans samples across all the markets were lower than the MRL of 0.01 mg kg⁻¹ set by the EU. Thus, all the above identified chemicals were found in brown beans samples from the study area but they are within the WHO MRL. However, in Ibadan and Ondo, Carbofuran was found to be above MRL. Pyrethroid residue found in white beans samples from the sampled markets are found to be within MRL.

3.1.3 Pyrethroids and Carbamates

The concentration of Amitraz and Cypermethrin I were lower than the EU's MRL in both brown and white beans (Table 1, Figure 2 and 3). Compared to EU's MRL of 0.02 mg kg⁻¹, white beans had higher levels of Flumioxazin across all markets except for Osun while Ado-Ekiti and Osun were the only ones that exceeded the EU MRL in the case of brown beans. Carbofuran was the only carbamate found in beans samples (Table 3, Figure 2 and 3). Its concentration was higher than the MRL stipulated by the EU in beans samples collected from Ondo as well as the brown beans from Ibadan and white beans from Osun. Pyrethroid residue found in white beans samples from the sampled markets are found to be within MRL.




Figure 2. Mean concentrations of pyrethroids and carbamates residues in brown beans samples. MRL₁ and MRL₂ are the maximum residue limit (mg kg⁻¹) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean.

All the above identified chemicals were found in brown beans samples from the study area but they are within the WHO MRL. However, in Ibadan and Ondo, Carbofuran was found to be above MRL.

3.2. Concentration of Identified Pesticide in Maize

3.2.1 Organochlorine

shows the mean concentrations of OCP residues in maize samples. δ -Lindane and β -Hexachlorocyclohexane were higher than the MRL of 0.01 mg kg⁻¹ set by the FAO/WHO and the EU in both yellow and white maize in all the markets. Likewise, the level of p,p'-DDT, Dieldrin were also higher than the MRL in white maize across all the markets and yellow maize in Ado-Ekiti, Ibadan and Ondo. While the concentrations of dieldrin in white maize of all the markets was higher than the MRL set by both the FAO/WHO and the EU, its concentration in yellow maize was lower than the FAO/WHO MRL but higher than the

Table 1

Mean concentrations (mg kg^{-1}) of organochlorine pesticide residues in bean samples obtained from selected markets in Southwest Nigeria.

Pesticide residue	MRL ₁	Concentration (mg kg^{-1})				
		MRL ₂	Ado-Ekiti	Ibadan	Osun	Ondo
Brown beans						
δ -Lindane	0.01	0.01	0.012 \pm 0.00	0.026 \pm 0.00	0.012 \pm 0.00	0.027 \pm 0.00
α -Lindane	0.01	0.01	0.032 \pm 0.00	0.032 \pm 0.00	0.027 \pm 0.00	0.033 \pm 0.00
β -Hexachlorocyclohexane	0.01	0.01	0.013 \pm 0.00	0.010 \pm 0.00	0.026 \pm 0.00	0.012 \pm 0.00
p,p'-DDT	0.1	0.05	0.200 \pm 0.01	0.200 \pm 0.02	0.230 \pm 0.00	0.249 \pm 0.01
Dieldrin	0.02	0.01	0.288 \pm 0.00	0.024 \pm 0.00	0.100 \pm 0.08	0.022 \pm 0.00
p,p'-DDE	0.1	0.05	0.190 \pm 0.00	0.209 \pm 0.00	0.173 \pm 0.00	0.187 \pm 0.01
Aldrin	0.02	0.01	0.012 \pm 0.00	0.011 \pm 0.00	0.010 \pm 0.00	0.013 \pm 0.00
White beans						
δ -Lindane	0.01	0.01	0.022 \pm 0.00	0.022 \pm 0.00	0.035 \pm 0.00	0.031 \pm 0.00
α -Lindane	0.01	0.01	0.036 \pm 0.00	0.030 \pm 0.00	0.034 \pm 0.00	0.033 \pm 0.00
β -Hexachlorocyclohexane	0.01	0.01	0.043 \pm 0.00	0.031 \pm 0.00	0.036 \pm 0.00	0.023 \pm 0.00
p,p'-DDT	0.1	0.05	0.217 \pm 0.00	0.211 \pm 0.00	0.213 \pm 0.00	0.214 \pm 0.00
Dieldrin	0.02	0.01	0.253 \pm 0.00	0.253 \pm 0.00	0.016 \pm 0.00	0.026 \pm 0.00
p,p'-DDE	0.1	0.05	0.196 \pm 0.00	0.213 \pm 0.01	0.349 \pm 0.00	0.210 \pm 0.01
Aldrin	0.02	0.01	0.012 \pm 0.00	0.013 \pm 0.00	0.017 \pm 0.00	0.011 \pm 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean \pm S.E. (n=3).

Table 2

Mean concentrations (mg kg^{-1}) of organophosphate pesticide residues in beans samples obtained from selected markets in Southwest Nigeria

Pesticide residue	MRL ₁	Concentration (mg kg^{-1})				
		MRL ₂	Ado-Ekiti	Ibadan	Osun	Ondo
Brown beans						
Phenthoate	N/A	N/A	0.023 \pm 0.00	0.034 \pm 0.00	0.014 \pm 0.00	0.033 \pm 0.00
Chlorthiophos	N/A	N/A	0.024 \pm 0.00	0.022 \pm 0.00	0.020 \pm 0.00	0.019 \pm 0.00
Ethion	N/A	0.01	0.014 \pm 0.00	0.012 \pm 0.00	0.012 \pm 0.00	0.010 \pm 0.00
Prothiofos	N/A	N/A	0.016 \pm 0.01	0.013 \pm 0.00	0.017 \pm 0.00	0.012 \pm 0.00
Iodofenphos	N/A	N/A	0.022 \pm 0.00	0.024 \pm 0.00	0.022 \pm 0.00	0.022 \pm 0.00
White beans						
Phenthoate	N/A	N/A	0.030 \pm 0.00	0.025 \pm 0.00	0.036 \pm 0.00	0.025 \pm 0.00
Chlorthiophos	N/A	N/A	0.024 \pm 0.00	0.021 \pm 0.00	0.042 \pm 0.00	0.014 \pm 0.00
Ethion	N/A	0.01	0.010 \pm 0.00	0.013 \pm 0.00	0.011 \pm 0.00	0.011 \pm 0.00
Prothiofos	N/A	N/A	0.023 \pm 0.00	0.045 \pm 0.00	0.035 \pm 0.00	0.011 \pm 0.00
Iodofenphos	N/A	N/A	0.022 \pm 0.00	0.023 \pm 0.00	1.230 \pm 0.01	0.027 \pm 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available. Values are expressed as mean \pm S.E. (n=3).

EU standard. On the bright side, the concentrations of Aldrin and Endosulfan in the maize samples across all the markets were lower than the MRL of 0.05 mg kg^{-1} set by the EU except for yellow maize samples collected from Ondo.

3.2.2 Organophosphates

The mean concentrations of the OPP residues are presented in Table 5. The concentration of Chlorthiophos, which was detected only in yellow maize, was between 0.018 and 0.049 mg kg^{-1} . The minimum and maximum values were observed in Osun and Ado-Ekiti, respectively. Similarly, Ethion was

not detected in all white and yellow maize samples as well as in yellow maize collected from Osun. Its concentrations in samples from Ondo was higher than the EU's MRL in samples obtained from Ondo. The levels of Malathion were higher than the FAO/WHO recommended MRL of 0.05 mg kg^{-1} in all samples of yellow and white maize obtained from Ado-Ekiti and Ibadan. The concentrations of Dichlorvos in both yellow and white maize samples were higher with MRL of 0.01 mg kg^{-1} set by the EU.

Conversely, the levels of pirimiphos methyl in both yellow and white maize samples were well below the MRL stipulated

Table 3

Mean concentrations (mg kg^{-1}) of pyrethroid and carbamate pesticide residues in bean samples obtained from selected markets in Southwest Nigeria.

Pesticide group	Pesticide residue	MRL ₁	Concentration (mg kg^{-1})				
			MRL ₂	Ado-Ekiti	Ibadan	Osun	Ondo
Brown beans Pyrethroid	Amitraz	N/A	0.00	0.016 ± 0.00	0.014 ± 0.00	0.012 ± 0.00	0.016 ± 0.00
	Flumioxazin	N/A	0.02	0.021 ± 0.00	0.020 ± 0.00	0.026 ± 0.00	0.016 ± 0.00
	Cypermethrin I	N/A	0.7	0.012 ± 0.00	0.015 ± 0.00	0.015 ± 0.00	0.014 ± 0.00
Carbamate	Carbofuran	N/A	0.00	0.003 ± 0.00	0.019 ± 0.00	0.006 ± 0.00	0.013 ± 0.00
White beans Pyrethroid	Amitraz	N/A	0.00	0.011 ± 0.00	0.014 ± 0.00	0.021 ± 0.00	0.015 ± 0.00
	Flumioxazin	N/A	0.02	0.021 ± 0.00	0.022 ± 0.00	0.017 ± 0.00	0.031 ± 0.00
	Cypermethrin I	N/A	0.7	0.010 ± 0.00	0.014 ± 0.00	0.015 ± 0.00	0.015 ± 0.00
	Carbamate	Carbofuran	N/A	0.00	0.010 ± 0.00	0.005 ± 0.00	0.025 ± 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available. Values are expressed as mean ± S.E. (n=3).




Figure 3. Mean concentrations of Pyrethroids and Carbamates residues in white beans samples. MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean (n=3).

by the FAO/WHO and slightly below that of the EU. Diazinon level in white maize were all higher than the MRLs while those of the yellow maize were all higher than the EU's MRL but generally lower than that of the FAO/WHO. Diazinon levels in maize collected from Osun had a mean value of 0.046 mg kg^{-1} which was very much higher than the stipulated MRLs. Chlorpyrifos was highest in yellow maize obtained from Ibadan but was not detected in Osun. In addition to Ibadan, its concentration in yellow maize from Ado-Ekiti was higher than MRLs. On the other hand, its concentrations in white maize across all markets were lower than the MRL set by FAO/WHO (0.05 mg kg^{-1}) but higher than that of the EU (0.01 mg kg^{-1}).

3.2.3 Pyrethroids and Carbamates

Amitraz was the only pyrethroid found in the maize sample (Table 6). Its concentration, which was in the range of $0.003 - 0.034 \text{ mg kg}^{-1}$, was lower than the EU MRL of 0.05 mg kg^{-1} . The concentration of Carbaryl in yellow maize obtained from Ibadan and Osun were higher than the MRLs values (Table 6). In the case of white maize, all samples across the markets recorded a range of $1.024 - 1.353 \text{ mg kg}^{-1}$, which were significantly higher than the 0.05 mg kg^{-1} and 0.02 mg kg^{-1} MRLs set by the FAO/WHO and EU, respectively. The concentration of Carbofuran in yellow maize for all the selected markets were lower than the MRLs (Figure 4). However, only Osun had a higher concentration of Carbofuran (0.206 mg kg^{-1}) when compared with FAO/WHO MRL of 0.01 mg kg^{-1} or EU's MRL of 0.05 mg kg^{-1} . On the other hand, white maize had a concentration of Carbofuran that was higher than the MRLs across all the four markets.

3.3. Concentration of Identified Pesticide in Millet

3.3.1 Organochlorine

The mean concentration of OCP residues in millet is presented in Table 7. More often, the levels of OCP residues in millet grains were higher than the MRL recommended by both the FAO/WHO as well as the EU. First, δ -Lindane, α -Lindane, β -Hexachlorocyclohexane and p,p'-DDT were higher than the MRLs. As for p,p'-DDE, Ado-Ekiti, Ibadan and Ondo all exceeded this limits substantially. While Dieldrin and Heptachlor levels in all millet samples ($0.040 - 0.127 \text{ mg kg}^{-1}$ and $0.020 - 0.046 \text{ mg kg}^{-1}$, respectively) were all higher

Table 4

Mean concentrations (mg kg^{-1}) of organochlorine pesticide residues in maize samples obtained from selected markets in Southwest Nigeria

Pesticide residue	Concentration (mg kg^{-1})				
	MRL ₁	MRL ₂	Adolbadan Ekiti	Osun	Ondo
Yellow maize					
δ -Lindane	0.01	0.01	0.025 ± 0.00	0.123 ± 0.00	0.021 ± 0.00
β -Hexachlorocyclohexane	0.01	0.01	1.000 ± 0.00	1.003 ± 0.00	0.999 ± 0.00
p,p'-DDT	0.1	0.05	2.170 ± 0.02	0.064 ± 0.00	2.157 ± 0.00
Dieldrin	0.02	0.01	0.013 ± 0.00	0.014 ± 0.00	0.014 ± 0.00
Aldrin	0.02	0.01	0.012 ± 0.00	0.012 ± 0.00	0.017 ± 0.00
Endosulfan	N/A	0.05	0.065 ± 0.00	0.018 ± 0.00	0.070 ± 0.05
White maize					
δ -Lindane	0.01	0.01	0.100 ± 0.00	0.982 ± 0.00	
β -Hexachlorocyclohexane	0.01	0.01	1.000 ± 0.00	0.026 ± 0.00	
p,p'-DDT	0.1	0.05	0.070 ± 0.00	0.019 ± 0.00	
Dieldrin	0.02	0.01	0.011 ± 0.00	0.021 ± 0.00	
Aldrin	0.02	0.01	0.016 ± 0.00	0.010 ± 0.00	
Endosulfan	N/A	0.05	0.020 ± 0.00	0.013 ± 0.00	

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available. Values are expressed as mean \pm S.E. (n=3). White maize was unavailable in Ondo market at the time of sample collection.

than EU's MRL of 0.01 mg kg^{-1} , none was over the 0.2 mg kg^{-1} set by the FAO/WHO. Aldrin followed a similar pattern but the 0.021 mg kg^{-1} concentration of millet obtained from Osun was higher than both MRLs. Chlordan was between the range of 0.007 and 0.035 mg kg^{-1} .

3.3.2 Organophosphates

The concentration of Phenthoate was in the range of $0.035 - 0.046 \text{ mg kg}^{-1}$ while Chlorthiophos and Prothiofos had ranges $0.022 - 0.035 \text{ mg kg}^{-1}$, and $0.017 - 0.024 \text{ mg kg}^{-1}$ (Table 8). The levels of Ethion in the millet grains from all the markets were above the MRL of 0.01 mg kg^{-1} set by the EU.

3.3.3 Pyrethroids

The concentration of Cypermethrin I in the millet samples was close and lower than the MRLs set by the FAO/WHO and the EU. On the other hand, millet samples obtained from Ado-Ekiti and Ibadan, which had a common mean concentration of Amitraz (0.040 mg kg^{-1}), were lower than the MRL of 0.05 mg kg^{-1} set by the EU while those of Osun and Ondo exceeded this limit.

3.3.4 Carbamates

The concentration of Carbofuran, the only identified carbamate in the millet grains, was higher than the MRL of 0.01 mg kg^{-1} set by the EU and lower than the 0.05 mg kg^{-1} set by

Table 5

Mean concentrations (mg kg^{-1}) of organophosphate pesticide residues in maize samples obtained from selected markets in Southwest Nigeria

Pesticide residue	MRL ₁	Concentration (mg kg^{-1})				
		MRL ₂	Ado-Ekiti	Ibadan	Osun	Ondo
Yellow maize						
Chlorthiophos	N/A	N/A	0.049 ± 0.00	0.044 ± 0.00	0.018 ± 0.00	0.044 ± 0.00
Ethion	N/A	0.0	0.008 ± 0.00	0.010 ± 0.00	N/D	0.052 ± 0.02
Malathion	0.05	N/A	0.840 ± 0.02	0.840 ± 0.02	0.397 ± 0.01	0.854 ± 0.02
Dichlorvos	N/A	0.0	1.110 ± 0.05	1.101 ± 0.02	0.985 ± 0.02	1.125 ± 0.00
Pirimiphos methyl	7	0.0	0.013 ± 0.00	0.018 ± 0.00	0.025 ± 0.00	0.015 ± 0.00
Diazinon	0.02	0.0	0.018 ± 0.00	0.016 ± 0.00	0.046 ± 0.00	0.015 ± 0.00
Chlorpyrifos	0.05	0.0	0.097 ± 0.07	0.216 ± 0.01	N/D	0.046 ± 0.00
White maize						
Malathion	0.05	N/A	0.381 ± 0.16	0.824 ± 0.00	0.035 ± 0.00	
Dichlorvos	N/A	0.0	1.099 ± 0.01	1.152 ± 0.02	0.113 ± 0.01	
Pirimiphos methyl	7	0.5	0.024 ± 0.00	0.044 ± 0.00	0.025 ± 0.00	
Diazinon	0.02	0.0	0.051 ± 0.00	0.037 ± 0.00	0.048 ± 0.00	
Chlorpyrifos	0.05	0.0	0.017 ± 0.00	0.029 ± 0.00	0.023 ± 0.00	

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available while "N/D" means undetected. Values are expressed as mean ± S.E. (n=3).

Table 6

Mean concentrations (mg kg^{-1}) of pyrethroid and carbamate pesticide residues in maize samples obtained from selected markets in Southwest Nigeria

Pesticide Group	Pesticide residue	MRL ₁	MRL ₂	Concentration (mg kg^{-1})			
				Ado-Ekiti	Ibadan	Osun	Ondo
Yellow maize							
Pyrethroid	Amitraz	N/A	0.05	0.005 ± 0.00	0.004 ± 0.00	0.023 ± 0.00	0.003 ± 0.00
Carbamate	Carbaryl	0.02	0.5	0.011 ± 0.00	0.202 ± 0.01	0.170 ± 0.00	0.013 ± 0.00
	Carbofuran	0.05	0.01	0.006 ± 0.00	0.006 ± 0.00	0.206 ± 0.00	0.005 ± 0.00
White maize							
Pyrethroid	Amitraz	N/A	0.05	0.034 ± 0.00	0.024 ± 0.00	0.032 ± 0.00	
Carbamate	Carbaryl	0.02	0.5	1.353 ± 0.02	1.107 ± 0.00	1.024 ± 0.00	
	Carbofuran	0.05	0.01	1.263 ± 0.03	0.036 ± 0.00	1.258 ± 0.00	

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available. Values are expressed as mean ± S.E. White maize was unavailable in Ondo market at the time of sample collection.

Table 7

Mean concentrations (mg kg^{-1}) of organochlorine pesticide residues in millet samples obtained from selected markets in Southwest Nigeria

Pesticide residue	MRL ₁	MRL ₂	Concentration (mg kg^{-1})			
			Ado-Ekiti	Ibadan	Osun	Ondo
δ -Lindane	0.01	0.01	0.015 ± 0.00	0.016 ± 0.00	0.011 ± 0.00	0.018 ± 0.00
α -Lindane	0.01	0.01	0.044 ± 0.00	0.044 ± 0.00	0.045 ± 0.00	0.146 ± 0.00
β -Hexachlorocyclohexane	0.01	0.01	0.025 ± 0.00	0.031 ± 0.00	0.032 ± 0.00	0.036 ± 0.00
p,p'-DDT	0.1	0.05	0.103 ± 0.00	0.103 ± 0.00	0.180 ± 0.01	0.037 ± 0.00
Dieldrin	0.2	0.01	0.040 ± 0.00	0.042 ± 0.00	0.046 ± 0.00	0.127 ± 0.00
p,p'-DDE	0.1	0.05	2.123 ± 0.01	2.107 ± 0.00	0.049 ± 0.00	1.127 ± 0.00
Heptachlor	0.2	0.01	0.046 ± 0.00	0.020 ± 0.00	0.036 ± 0.00	0.040 ± 0.00
Chlordan	N/A	N/A	0.007 ± 0.00	0.012 ± 0.00	0.009 ± 0.00	0.035 ± 0.03
Aldrin	0.2	0.01	0.012 ± 0.00	0.011 ± 0.00	0.021 ± 0.00	0.012 ± 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available. Values are expressed as mean ± S.E. (n=3).

Table 8

Mean concentrations (mg kg^{-1}) of organophosphate, pyrethroid and carbamate pesticide residues in millet samples obtained from selected markets in Southwest Nigeria.

Pesticide group	Pesticide residue	Concentration (mg kg^{-1})					
		MRL ₁	MRL ₂	Ado-Ekiti	Ibadan	Osun	Ondo
Organophosphate	Phenthoate	N/A	N/A	0.046 ± 0.00	0.041 ± 0.00	0.035 ± 0.00	0.046 ± 0.00
	Chlorthiophos	N/A	N/A	0.033 ± 0.00	0.022 ± 0.00	0.034 ± 0.00	0.035 ± 0.00
	Ethion	N/A	0.01	0.022 ± 0.00	0.013 ± 0.00	0.014 ± 0.00	0.013 ± 0.00
	Prothiofos	N/A	N/A	0.017 ± 0.00	0.022 ± 0.00	0.024 ± 0.00	0.022 ± 0.00
Pyrethroid	Cypermethrin I	0.3	0.7	0.016 ± 0.00	0.016 ± 0.00	0.015 ± 0.00	0.015 ± 0.00
	Amitraz	N/A	0.05	0.040 ± 0.00	0.040 ± 0.00	0.193 ± 0.00	0.071 ± 0.03
Carbamates	Carbofuran	0.05	0.01	0.036 ± 0.00	0.032 ± 0.00	0.017 ± 0.00	0.027 ± 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. "N/A" means not available. Values are expressed as mean ± S.E. Different alphabets in the same row mean significant difference at $P < 0.05$.




Figure 4. Mean concentrations of pyrethroids and carbamates residues in yellow maize samples, MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean ($n=3$).

FAO/WHO across all four markets (Table 8).

3.4. Concentration of Identified Pesticide in Rice

3.4.1 Organochlorine

The concentration of OCP residues in rice samples is given in Table 9. δ -Lindane, α -Lindane, exceeded the MRL of 0.01 mg kg^{-1} set by both the FAO/WHO and the EU. β -Hexachlorocyclohexane showed a similar trend as rice samples from all market exceeded the MRL except for rice grains collected in Osun. Meanwhile, the concentration of p,p'-DDT and p,p'-DDE in the rice grains were lower than the MRLs in all the markets. Lastly, Aldrin level in rice grains obtained from Ibadan and Osun were lower than the MRL of 0.01 mg kg^{-1} set by the EU while those of Ondo and Ado-Ekiti were higher than this limit but lower than the 0.02 mg kg^{-1} set by the FAO/WHO. The concentration of Dieldrin in any one of the markets was higher than at least one of the MRLs.

3.4.2 Organophosphates, Pyrethroids and Carbamates

Phenthoate, Chlorthiophos, Ethion, Iodophenos and Prothiofos concentrations in the rice grains were in the range of $0.023 - 0.045 \text{ mg kg}^{-1}$, $0.005 - 0.026 \text{ mg kg}^{-1}$, $1.13 - 1.460 \text{ mg kg}^{-1}$, $0.005 - 0.007 \text{ mg kg}^{-1}$, $1.50 - 1.528 \text{ mg kg}^{-1}$, respectively (Table 10 & Figure 5). The minimum mean

level of Ethion was at least 100 times higher than the MRL of 0.01 mg kg^{-1} set by the EU. Table 10 : Mean concentrations (mg kg^{-1}) of organophosphate, pyrethroid and carbamate pesticide residues in rice samples obtained from selected markets in Southwest Nigeria. The concentrations of Cypermethrin I in rice was lower than the MRL recommended by the EU (Table 10). Also, Flumioxazin was lower than the MRL of 0.02 mg kg^{-1} set by the EU, except for the rice samples obtained from Ondo. However, Amitraz was generally higher than the set limit of 0.05 mg kg^{-1} set by the FAO. Only carbofuran was detected in the rice grain samples and its concentration was above the MRL of 0.01 mg kg^{-1} set by the EU or FAO across all markets (Table 10).




Figure 5. Mean concentrations of organophosphate, pyrethroids and carbamate pesticide residues in rice samples. MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean ($n=3$).

Ethion, Prothiofos and amthraz are dominantly present but below WHO and FAO MRL in grain samples from South-west Nigeria.

4. DISCUSSION

4.1. Determine the Concentration of identified pesticides in grains

Rachel Carson, "a marine biologist and author, is widely regarded as the founder of the global environmental movement. Carson advocated for environmental protection and conservation, but she also granted numerous warnings about

Table 9

Mean concentrations (mg kg^{-1}) of organochlorine pesticide residues in rice samples obtained from selected markets in Southwest Nigeria

Pesticide residue	MRL ₁	Concentration (mg kg^{-1})				
		MRA ₂	Ondo-Ekiti	Ibadan	Osun	Ondo
δ -Lindane	0.01	0.01	0.033 \pm 0.00	0.035 \pm 0.00	0.040 \pm 0.00	0.035 \pm 0.00
α -Lindane	0.01	0.01	0.031 \pm 0.00	1.032 \pm 0.00	1.010 \pm 0.00	0.699 \pm 0.33
β -Hexachlorocyclohexane	0.01	0.01	0.018 \pm 0.00	0.019 \pm 0.00	0.006 \pm 0.00	0.020 \pm 0.00
p,p'-DDT	0.1	0.05	0.010 \pm 0.00	0.016 \pm 0.00	0.015 \pm 0.00	0.011 \pm 0.00
Dieldrin	0.02	0.01	0.012 \pm 0.00	0.025 \pm 0.00	2.112 \pm 0.00	0.015 \pm 0.00
p,p'-DDE	0.1	0.05	0.021 \pm 0.00	0.016 \pm 0.00	0.013 \pm 0.00	0.014 \pm 0.00
Aldrin	0.02	0.01	0.017 \pm 0.01	0.006 \pm 0.00	0.004 \pm 0.00	0.013 \pm 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean \pm S.E. (n=3).

Table 10

Mean concentrations (mg kg^{-1}) of organophosphate, pyrethroids and carbamate pesticide residues in rice samples

Pesticide group	Pesticide residue	MRL ₁	Concentration (mg kg^{-1})				
			MRA ₂	Ondo-Ekiti	Ibadan	Osun	Ondo
Organophosphate	Phenthoate	N/A	N/A	0.023 \pm 0.00	0.045 \pm 0.00	0.027 \pm 0.00	0.027 \pm 0.00
	Chlorthiophos	N/A	N/A	0.005 \pm 0.00	0.007 \pm 0.00	0.008 \pm 0.00	0.026 \pm 0.00
	Ethion	N/A	0.01	0.450 \pm 0.00	1.460 \pm 0.02	1.315 \pm 0.00	1.130 \pm 0.02
	Iodofenphos	N/A	N/A	0.007 \pm 0.00	0.005 \pm 0.00	0.007 \pm 0.00	0.009 \pm 0.00
	Prothiofos	N/A	N/A	0.502 \pm 0.01	1.508 \pm 0.00	1.500 \pm 0.01	1.528 \pm 0.02
Pyrethroid	Cypermethrin I	2	2	0.018 \pm 0.00	0.003 \pm 0.00	0.002 \pm 0.00	0.005 \pm 0.00
	Flumioxazin	N/A	0.02	0.011 \pm 0.00	0.014 \pm 0.00	0.011 \pm 0.00	0.022 \pm 0.00
	Amitraz	N/A	0.05	0.300 \pm 0.03	1.449 \pm 0.01	0.007 \pm 0.00	1.337 \pm 0.01
Carbamate	Carbofuran	N/A	0.01	0.016 \pm 0.00	0.017 \pm 0.00	0.017 \pm 0.00	0.017 \pm 0.00

MRL₁ and MRL₂ are the maximum residue limit (mg kg^{-1}) set by the FAO/WHO and the European Union, respectively. Values are expressed as mean \pm S.E. (n=3).

pesticides' dangers to humans and the environment (Bonzi, 2013). The use of DDT (dichlorodiphenyltrichloroethane), an organochlorine insecticide that was critical during WWII in combating typhus, yellow fever, as well as malaria-transmitting insects, was at the heart of this debate" (D'Amato et al., 2002). DDT was used in every malaria control effort between 1946 and 1970 because of how well it worked to eliminate mosquitos that spread the disease. Notwithstanding, due to its high toxicity, the use of this insecticide was reduced in later years. The Stockholm Convention on Persistent Organic Pollutants emphasizes the need for DDT alternatives by citing its environmental persistence and bioaccumulation (UNEP, 2008; Vanden Bilcke, 2002). Pesticide residues have been found to disperse across the environment in recent years, polluting several ecosystems as well as threatening food and water supplies. Because growth in population might be inconceivable without a rise in agricultural output, which is directly related to the use of pesticides and fertilizers, this contamination is observed to be a result of population growth (Asiegbu et al., 2022; Carvalho, 2017; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Pesticides are a broad category of organic compounds with a wide range of physicochemical properties. They are used to prevent and control pests in

a variety of crops and plantations, increasing productivity, but have a growing number of negative environmental and human health effects that are often caused by agrochemical changes. The need for notable changes in the global food system has been spotlighted in multiple reports. Agriculture should indeed confront the dual hurdles of feeding an expanding population even while expanding its global environmental impact. Even though pesticides were originally intended to improve agricultural output and prevent the spread of infectious diseases, their negative consequences have now outweighed those benefits. Pesticides have had such an impact on our ecosystem because of their persistent nature that they have infiltrated many food chains and higher trophic levels, including those of humans and other large mammals (M.O. Raimi, 2019; M.O. Raimi et al., 2020; M.O. Raimi, Olawale, et al., 2022). Contaminated water (Afolabi & Morufu, 2021; Afolabi & Raimi, 2021; Mo & Sabinus, 2017; O.R. Morufu, Clinton, & Bowale, 2021; O.R. Morufu, Henry, et al., 2021; O.R. Morufu, Olawale, & Clinton, 2021; R. Morufu & E, 2017; Odipe et al., 2018; A.S. Olalekan et al., 2022; M.R. Olalekan, Albert, et al., 2022; M.R. Olalekan, Olawale, et al., 2022; R.M. Olalekan et al., 2018; Omotoso et al., 2021; M. Raimi & Sawyerr, 2022; M.O. Raimi et al., 2021; M.O. Raimi, Sawyerr, et al., 2022; O. Raimi et al., 2022; Stephen et al., 2022), air (Ajayi et al., 2020; Ic et al., 2022; M.O. Raimi et al., 2018), or food (Adiama

et al., 2022; Habeeb et al., 2022; H.M. Lateefat et al., 2022; M.H. Lateefat, Opasola, Adiamata, et al., 2022; M.H. Lateefat, Opasola, Misbahu, & Morufu, 2022) consumption has now been interconnected to a variety of acute and chronic human disorders. Numerous pesticides have been linked to health and environmental problems (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; M.R. Olalekan et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020), and several pesticides are no longer used in agriculture. Pesticides can be ingested, inhaled, or applied directly to the skin. The type of pesticide, the length of exposure, the method of exposure, and the individual health status (such as dietary deficits and the condition of one's skin, for example) all affect the potential health outcome. Pesticides may be metabolized, expelled, stored, or bioaccumulated in body fat within a human or animal body. Chemical pesticides have been linked to a wide range of unfavorable health effects, including impacts on the skin, gastrointestinal system, nervous system, respiratory system, reproductive system, and endocrine system (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; M.R. Olalekan et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020). Additionally, excessive levels of occupational, unintentional, or purposeful pesticide exposure can cause hospitalization and even death (Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020). Pesticide residues have been found in a variety of common foods and beverages, including prepared meals, water, wine, fruit juices, snacks, and animal feeds. Besides this, it should be understood that washing as well as peeling will not completely remove residues. The majority of the time, the concentrations are lower than the legal safe levels. These "safe limits" could, even so, underestimate the true health risk, as in the case of simultaneous exposure to two or more chemicals, which occurs in real-life situations and may have synergistic effects. Thus, this study revealed concentration levels of organochloride pesticides and organo-phosphate pesticides in grain samples drawn from selected markets in southwest Nigeria. The nutrient composition of cereals is an important indicator of its quality and the overall health benefits of its consumption. First, we observed large variations in moisture content of the different grain samples. These differences can be linked to factors such as planting methods and management practices, processing, environmental factors, pests and diseases etc. (Stathers et al., 2020; Zhang et al., 2021). The ash content of grains depends very much on their mineral composition (Babarinde et al., n.d.). The ash content of legumes has been generally higher than those of cereals (??). Similarly, higher ash content in beans was higher than those of the cereals in our study. The protein composition was the highest in cowpea and it fell within the range of 20 – 30% reported by Odogwu et al. (2021). Also, cowpeas was around 11% reported by ?. This is because the removal of the bran and germ of cereals leaves

it with the endosperm which lacks nutrients but highly rich in carbohydrate (?). Pigmented maize was recommended by ? due to its high nutritional value and presence of antioxidants, hence its high protein composition, as against white maize, as observed in our study. However, the higher ash content, crude fibre and crude fat in white maize we observed may be attributed to varietal factors and performance (?). It is well known that rice has a higher composition of carbohydrate due to its high starch content while maize and millet have better crude protein and fat (?). In our study as rice also had the highest composition of carbohydrate and which ranged between 76.92 and 86.03% similar to that reported by ? who compared several varieties of rice in Nigeria. Meanwhile, maize and millet had a higher crude protein and fat level s. Pesticides are commonly applied in preserving food products in Nigeria. Hence, they have been widely reported in food crops (Ajayi et al., 2020; Asiegbu et al., 2022; Hussain et al., 2021a, 2021b; Isah, Raimi, et al., 2020; Isah, Sawyerr, et al., 2020; O.R. Morufu, 2021; O.R. Morufu, Tonye, et al., 2021; R.M. Olalekan et al., 2020; M.O. Raimi et al., 2020; Sosan et al., 2018). Some OCPs in this study such as β -HCH, Heptachlor, Endosulfan, Aldrin, Lindane, Dieldrin, p,p'-DDE, p,p'-DDT have been previously reported in maize, beans, millet and rice in Nigeria (????). Similarly, (?) reported Dichlorvos, Carbofuran, Carbaryl, Diazinon, Pirimiphos-methyl, Malathion, Chlorpyrifos as the OPPs and carbamates in beans, rice, wheat and maize samples collected from Ondo State. Additional pesticide residues such as those of Phenthoate and Iodofenphos found in our study could be as a result of the four (4) different locations and larger sample size. Amitraz, Cypermethrin and Flumioxazin, which were the pyrethroids residues observed in our study had been reported in grain samples in Nigeria (?). This suggests the widespread use of various groups of pesticides including organochlorine, organophosphate, carbamates and pyrethroids for grain preservation across Nigeria. In cowpea and maize grains obtained directly from farms in Ghana, ? reported the mean concentration of organochlorine, organophosphates, and pyrethroids ranging from 0.001 to 0.103 mg kg⁻¹, 0.002 to 0.019 mg kg⁻¹ and 0.002 to 0.028 mg kg⁻¹ in maize, and 0.001 to 0.108 mg kg⁻¹, 0.002 to 0.015 mg kg⁻¹, 0.001 to 0.039 mg kg⁻¹ in cowpea, respectively. Our values were generally higher than the upper limits of these range of values because higher doses of pesticides must have been applied during postharvest storage. The concentration of OCP residues in maize sample in this study were comparable to those of ? in Nassarawa State which reported a range of 0.03 to 0.13 mg/kg, 0.018 to 0.02 mg/kg, 0.005 to 0.02 mg/kg, 0.25 to 1.25 mg/kg, and 0.04 mg/kg for Aldrin, Dieldrin, Endosulfan, Lindane, and DDT, in maize and guinea corn respectively. While Aldrin and Dieldrin was within these range of values, the upper limit of Endosulfan, Lindane and DDT were much higher in our study. The results of ? showed that Dieldrin, and Endosulfan while, p,p'-DDT and p,p'-DDE concentrations were in the range of 0.99 – 11.01 mg/kg, 0.85 – 1.37 mg/kg, 1.05 – 1.82 mg/kg and 0.48 – 1.23 mg/kg, respectively in beans from markets in Ibadan metropolis. Meanwhile, in our study, Endosulfan was

undetected, Dieldrin was within this range, p,p'-DDT and p,p'-DDE were much lower. Variations of such nature have been widely reported because the concentrations of pesticide residues in grains depend on several factors such as merchant attitude and knowledge, type and quantity of pesticide used for preservation, duration of exposure of the grains etc.

5. CONCLUSION

Pesticides have negative effects on society, so it is important to reevaluate their viability. To meet the need for food globally, alternative routes must be pursued to provide healthier food. Therefore, one of the primary findings of this study is that contamination is universal because no individual of any species throughout the entire basin appeared to be free of numerous pesticide residues in grains. The results frequently showed high inter- and intra-specific contamination, which makes sense given the target market and dietary diversity in the area. Therefore, efforts to locate and remove pollution sources will have an impact throughout Southwest Nigeria, not just on the markets that are previously recognized to be the most polluted. This study found that grains purchased from particular markets in southwest Nigeria contained numerous pesticide residues. The MRLs set by the EU or FAO/WHO or both were surpassed by 17 out of the total 27 pesticides reported in this work in at least one grain, despite the fact that there were no published codex MRLs for some pesticide residues in some grains. These residues were classified among the four classes of pesticides: carbamates, organochlorines, organophosphates, and pyrethroids. In actuality, 90% of the mainly banned organochlorine pesticides exceeded MRLs. The merchants who typically admit to the usage of pesticides were connected to their misuse; almost half of these merchants don't even read the instructions on pesticide containers before using them. There is no doubt that the populous is exposed to the dangers brought on by eating wheat polluted with pesticides. While, cancer and physiological abnormalities are among the hazards. According to this research, the sale of beans, maize, millet, and rice grains in some markets in Southwest Nigeria may pose a serious health risk to the general public. In conclusion, routine assessments will give information on the contamination level in real time as well as the potential effects of unforeseen occurrences or subsequent actions controlling the flow of pollution. The sale of pesticides must be controlled in order to ensure that agricultural workers are trained in the safe use of these chemicals, and studies into the long-term effects of pesticide use on agricultural workers and the children who live in their homes are crucial as the volume of pesticide imports keeps rising. The Ministry of Health and Ministry of Agriculture employees can better target their educational efforts with the support of the information given above.

CONFLICTS OF INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

ACKNOWLEDGEMENTS

We note with sadness the passing of our friend, colleague, and mentor Prof. Mynepalli K. C. Sridhar; working with him was wonderful and we strongly believe that he would have loved to coauthor this research paper – we miss him. We would like to say special thanks to all anonymous reviewer for providing very helpful comments for their valuable suggestions for improving this manuscript.

ORCID

Modupe Abeke Oshatunberu [0000-0002-8886-6575](https://orcid.org/0000-0002-8886-6575)
 Adebayo Oladimeji [0000-0002-3307-5047](https://orcid.org/0000-0002-3307-5047)
 Sawyerr Olawale Henry [0000-0003-3897-9034](https://orcid.org/0000-0003-3897-9034)
 Opasola Afolabi Olaniyan [0000-0002-4962-5744](https://orcid.org/0000-0002-4962-5744)
 Morufu Olalekan Raimi [0000-0001-5042-6729](https://orcid.org/0000-0001-5042-6729)

A. SUPPLEMENTARY INFORMATION

Supplementary information to this article can be found online at <https://doi.org/10.53365/nrhh/171368>.

FUNDING

Not Applicable (No funds were granted, received or used).

ETHICAL APPROVAL

This study entitled: “Evaluation of Pesticide Residues in Grain Sold at Selected Markets of Southwest Nigeria”, which was submitted for ethical approval on August 15, 2020 to the Research Ethics Committee of Kwara State University. For the development of this study, the fundamental ethical principles for research involving human beings, described and established by the Resolution 466/2012 and its complementary ones, was considered and approval was granted by the Kwara State University research ethics committee.

During the course of this research work, the participants were accorded the due respect so as to ensure co-operation and information collected were treated with utmost confidentiality. The cultures of the community were also respected during the course of the research work. Informed consent was obtained from all of the participants.

AUTHOR CONTRIBUTIONS

Modupe Abeke Oshatunberu: Conceptualization, Investigation, Writing - original draft, Writing - review & editing. Modupe Abeke Oshatunberu and Adebayo Oladimeji: Investigation, Writing - original draft, Writing - review & editing. Adebayo Oladimeji and Sawyerr Olawale Henry: Resources, supervising, review & editing. Modupe Abeke Oshatunberu, Morufu Olalekan Raimi: Methodology, Project administration, Writing - original draft, Writing - review & editing. Morufu Olalekan Raimi: Data curation, Formal analysis, Writing - original draft, Writing - review & editing. Modupe Abeke Oshatunberu, Adebayo Oladimeji, Sawyerr Olawale Henry:

Conceptualization, Data curation, Funding acquisition, Project administration, Writing - original draft, Writing - review & editing.

REFERENCES

- Adiama, Y.B., Sawyerr, O.H., Olaniyi, O.A., Fregene, A.F., Alabede, M., Raimi, Mo., 2022. Assessment of Microbiological Quality of Ready to Eat Food Served in Ships Along Warri, Koko and Port Harcourt Water Ways, Nigeria. *Online Journal of Microbiological Research*. 1(1), 1–7. <https://doi.org/10.31586/ojmr.2021.230>
- Afolabi, A.S., Morufu, O.R., 2021. Investigating Source Identification and Quality of Drinking Water in Piwoyi Community of Federal Capital Territory, Abuja Nigeria. <https://doi.org/10.21203/rs.3.rs-736140/v1>
- Afolabi, A.S., Raimi, M.O., 2021. When Water Turns Deadly: Investigating Source Identification and Quality of Drinking Water in Piwoyi Community of Federal Capital Territory. *Online Journal of Chemistry*. 1, 38–58. <https://doi.org/10.31586/ojc.2021.010105>
- Ajayi, F.A., Raimi, M.O., Steve-Awogbami, O.C., Adeniji, A.O., Adebayo, P.A., 2020. Policy Responses to Addressing the Issues of Environmental Health Impacts of Charcoal Factory in Nigeria: Necessity Today. *Essentiality Tomorrow. Communication, Society and Media*. 3(3). <https://doi.org/10.22158/csm.v3n3p1>
- AOAC., 2000. *Official Methods of Analysis*. Association of Official Analytical Chemists, Washington DC.
- Asiegbu, O.V., Ezekwe, I.C., Mo, R., 2022. Assessing pesticides residue in water and fish and its health implications in the Ivo river basin of South-eastern Nigeria. *MOJ Public Health*. 11(4), 136–142. <https://doi.org/10.15406/mojph.2022.11.00390>
- Babarinde, G.O., Adeyanju, J.A., Ogunleye, K.Y., Adegbola, G.M., Eburn, A.A., Wade, D., n.d.
- Bonzi, R.S., 2013. Meio século de Primavera silenciosa: um livro que mudou o mundo. *Desenvolv. e Meio Ambient. Ed. UFPR*. 28, 207–215. <https://doi.org/10.5380/dma.v28i0.31007>
- Carriger, J.F., Rand, G.M., Gardinali, P.R., Perry, W.B., Tompkins, M.S., Fernandez, A.M., 2006. Pesticides of potential ecological concern in sediment from South Florida canals: an ecological risk prioritization for aquatic arthropods. *Soil and Sediment Contamination*. 15, 21–45. <https://doi.org/10.1080/15320380500363095>
- Carvalho, F.P., 2017. Pesticides, environment, and food safety. *Food and Energy Security*. 6, 48–60. <https://doi.org/10.1002/fes3.108>
- D'amato, C., Torres, J.P.M., Malm, O., 2002. DDT (Dicloro difenil tricloretoano): toxicidade e contaminação ambiental - Uma revisão *Quim. Nova*. 25. <https://doi.org/10.1590/S0100-40422002000600017>
- Ekpa, O., Palacios-Rojas, N., Kruseman, G., Fogliano, V., Linne-mann, A.R., 2018. Sub-Saharan African maize-based foods: Technological perspectives to increase the food and nutrition security impacts of maize breeding programmes. *Global Food Security*. 17, 48–56. <https://doi.org/10.1016/j.gfs.2018.03.007>
- Habeeb, M.L., Opasola, A.O., Garba, M., Olalekan, M.R., 2022. A Wake-Up Call: Determination of Antibiotics Residue Level in Raw Meat in Abattoir and Selected Slaughterhouses in Five Local Government in Kano State, Nigeria. *Journal of Veterinary Health Science*. 3(1), 54–61. <https://doi.org/10.33140/JVHS.03.01.02>
- Hussain, M.I., Morufu, O.R., Henry, O.S., 2021a. Patterns of Chemical Pesticide Use and Determinants of Self-Reported Symptoms on Farmers Health: A Case Study in Kano State for Kura Local Government Area of Nigeria. *Research on World Agricultural Economy*. 2(1), 37–48. <https://doi.org/10.36956/rwae.v2i1.342>
- Hussain, M.I., Morufu, O.R., Henry, O.S., 2021b. Probabilistic Assessment of Self-Reported Symptoms on Farmers Health: A Case Study in Kano State for Kura Local Government Area of Nigeria. *Research on World Agricultural Economy*. 2(1), 1–11. <http://dx.doi.org/10.36956/rwae.v2i1.336>
- Ic, C.-E., Ic, O., Ezekwe, I.C., Mo, R., 2022. Slow death from pollution: potential health hazards from air quality in the mgbede oil fields of south-south Nigeria. *Open Access Journal of Science*. 5(1), 61–69. <https://doi.org/10.15406/oajs.2022.05.00177>
- ILNAS-EN., 2018. <https://ilnas.services-publics.lu/ecnor/downloadP>
- Isah, H.M., Raimi, M.O., Sawyerr, H.O., Odipe, O.E., Bashir, B.G., Suleiman, H., 2020. Qualitative Adverse Health Experience Associated with Pesticides Usage among Farmers from Kura. *Merit Research Journal of Medicine and Medical Sciences*. 8(8), 432–447. <https://doi.org/10.5281/zenodo.4008681>
- Isah, H.M., Sawyerr, H.O., Raimi, M.O., Bashir, B.G., Haladu, S., Odipe, O.E., 2020. Assessment of Commonly Used Pesticides and Frequency of Self-Reported Symptoms on Farmers Health in Kura, HolyKnight. *JELM*, pp. 31–54. <http://dx.doi.org/10.46410/jelm.2020.1.1.05>
- James, S., Nwabueze, T.U., Onwuka, G.I., Ndife, J., Usman, M.A., 2020. Chemical and nutritional composition of some selected lesser known legumes indigenous to Nigeria. *Heliyon*. 6(11), e05497. <https://doi.org/10.1016/j.heliyon.2020.e05497>
- Lateefat, H.M., Faith, A., Yusuf, A.B., Raimi, O.M., 2022. Food for the Stomach Nourishing our Future: Assessment of Potassium Bromate in Local and Packaged Bread Sold in Ilorin Metropolis. *Public Health Open Access*. 6(1), 000197. <https://doi.org/10.23880/phoa-16000197>
- Lateefat, M.H., Opasola, A.O., Adiama, B.Y., Ibrahim, A., Morufu, O.R., 2022. Elixirs of Life, threats to Human and Environmental Well-being: Assessment of Antibiotic Residues in Raw Meat Sold Within Central Market Kaduna Metropolis, Kaduna State, Nigeria. *bioRxiv*. <https://doi.org/10.1101/2022.01.04.474997>
- Lateefat, M.H., Opasola, A.O., Misbahu, G., Morufu, O.R., 2022. A Wake-Up Call: Determination of Antibiotics Residue Level in Raw Meat in Abattoir and Selected Slaughterhouses in Five Local Government in Kano State, Nigeria. *bioRxiv*. <https://doi.org/10.1101/2022.01.04.474991>
- Mo, R., Sabinus, C.E., 2017. An Assessment of Trace Elements in Surface and Ground Water Quality in the Ebocha-Obrikom Oil and Gas Producing Area of Rivers State, Nigeria. *International Journal for Scientific and Engineering Research*. 8(6), 2229–5518.
- Morufu, O.R., 2021. Self-reported Symptoms on Farmers Health and Commonly Used Pesticides Related to Exposure in Kura. *Annals of Community Medicine & Public Health*. 1(1), 1002–1002.
- Morufu, O.R., Clinton, I.E., Bowale, A., 2021. Statistical and Multivariate Techniques to Trace the Sources of Ground Water Contaminants and Affecting Factors of Groundwater Pollution in an Oil and Gas Producing Wetland in Rivers State, Nigeria. *medRxiv*. <https://doi.org/10.1101/2021.12.26.21268415>
- Morufu, O.R., Henry, O.S., Clinton, I.E., Gabriel, S., 2021. Many Oil Wells, One Evil: Potentially toxic metals concentration, seasonal variation and Human Health Risk Assessment in Drinking Water Quality in Ebocha-Obrikom Oil and Gas Area of Rivers State, Nigeria. <https://doi.org/10.1101/2021.11.06.21266005>
- Morufu, O.R., Olawale, H.S., Clinton, I.E., 2021. Quality water not everywhere: Exploratory Analysis of Water Quality Across Ebocha-Obrikom Oil and Gas Flaring Area in the Core Niger Delta Region of Nigeria. PREPRINT (Version 1) available at Research Square. <https://doi.org/10.21203/rs.3.rs-953146/v1>
- Morufu, O.R., Tonye, V.O., Ogah, A., Henry, A.E., Abinotami, W.E.,

2021. Articulating the effect of Pesticides Use and Sustainable Development Goals (SDGs): The Science of Improving Lives through Decision Impacts. *Research on World Agricultural Economy*. 2(1), 29–36. <https://doi.org/10.36956/rwae.v2i1.347>
- Morufu, R., E, C., 2017. Assessment of Trace Elements in Surface and Ground Water Quality, and others, (Eds.). LAP Lambert Academic Publishing, Mauritius, pp. 978–981.
- Mostafalou, S., Abdollahi, M., 2012. Concerns of environmental persistence of pesticides and human chronic diseases. *Clinical and Experimental Pharmacology*. S5, e002. <https://doi.org/10.4172/2161-1459.S5-e002>
- Odipe, O.E., Raimi, M.O., Suleiman, F., 2018. Assessment of Heavy Metals in Effluent Water Discharges from Textile Industry and River Water at Close Proximity: A Comparison of Two Textile Industries from Funtua and Zaria. *Madridge Journal of Agriculture and Environmental Sciences*. 1(1), 1–6. <https://doi.org/10.18689/mjaes-1000101>
- Odogwu, B.A., Uzogara, M.O., Worlu, H., Agbagwa, Io., 2021. Factors affecting stakeholders' preferences for cowpea grains in selected parts of Nigeria. *Agriculture, Nutrition and Development*. 21(3), 17669–17681. <https://doi.org/10.18697/ajfand.98.19890>
- Ogisi, O.D., Begho, T., 2021. Covid 19: Ramifications for progress towards the sustainable development goals (SDGs) in Nigeria. *International Review of Applied Economics*. 35(2), 256–268. <https://doi.org/10.1080/02692171.2020.1864302>
- Olalekan, A.S., Adewoye, S.O., Henry, S.O., 2022. Comparative Assessment of Seasonal Variations in the Quality of Surface water and its associated health hazards in Gold Mining Areas of Osun State, South-West Nigeria. PREPRINT (Version 1) available at Research Square. <https://doi.org/10.21203/rs.3.rs-2245715/v1>
- Olalekan, M.R., Abiola, I., Ogah, A., Dodeye, E.O., 2021. Exploring How Human Activities Disturb the Balance of Biogeochemical Cycles: Evidence from the Carbon, Nitrogen and Hydrologic Cycles. *Research on World Agricultural Economy*. 2, 23–44. <https://doi.org/10.36956/rwae.v2i3.426>
- Olalekan, M.R., Albert, O., Iyingiala, A.A., Sanchez, D.N., Telu, M., 2022. An environmental/scientific report into the crude oil spillage incidence in Tein community. *Journal of Environmental Chemistry and Toxicology*. 6(4), 1–6.
- Olalekan, M.R., Olawale, H.S., Ie, C., Oposola, A.O., 2022. Quality Water, Not Everywhere: Assessing the Hydrogeochemistry of Water Quality across Ebocha-Obrikom Oil and Gas Flaring Area in the Core Niger Delta Region of Nigeria. *Pollution*. 8, 751–778.
- Olalekan, R.M., Muhammad, I.H., Okoronkwo, U.L., Akopjubaro, E.H., 2020. Assessment of safety practices and farmer's behaviors adopted when handling pesticides in rural Kano state, Nigeria. *Arts & Humanities Open Access Journal*. 4(5), 191–201. <https://doi.org/10.15406/ahoaj.2020.04.00170>
- Olalekan, R.M., Omidiji, A.O., Nimisingha, D., Odipe, O.E., Olalekan, A.S., 2018. Health Risk Assessment on Heavy Metals Ingestion through Groundwater Drinking Pathway for Residents in an Oil and Gas Producing Area of Rivers State, Nigeria. *Open Journal of Yangtze Gas and Oil*. 3, 191–206. <https://doi.org/10.4236/ojogas.2018.33017>
- Olalekan, R.M., Omidiji, A.O., Williams, E.A., Christianah, M.B., Modupe, O., 2019. The roles of all tiers of government and development partners in environmental conservation of natural resource: a case study in Nigeria. *MOJ Ecology & Environmental Sciences*. 4(3), 114–121. <https://doi.org/10.15406/mojes.2019.04.00142>
- Omotoso, A.J., Omotoso, E.A., Morufu, O.R., 2021. Potential Toxic levels of Cyanide and Heavy Metals in Cassava Flour Sold in Selected Markets in Oke Ogun Community, Oyo State, Nigeria. PREPRINT (Version 1) available at Research Square. <https://doi.org/10.21203/rs.3.rs-658748/v1>
- Pan-Germany., 2012. www.pangermany.org/download/Vergift_EN-201112-web.pdf
- Raimi, M., Sawyerr, H., 2022. Preliminary Study of Groundwater Quality Using Hierarchical Classification Approaches for Contaminated Sites in Indigenous Communities Associated with Crude Oil Exploration Facilities in Rivers State Nigeria. *Open Journal of Yangtze Oil and Gas*. 7, 124–148. <https://doi.org/10.4236/ojogas.2022.72008>
- Raimi, M.O., 2019. 21st Century Emerging Issues in Pollution Control. 6th Global Summit and Expo on Pollution Control. .
- Raimi, M.O., Abiola, O.S., Atoyebi, B., Okon, G.O., Popoola, A.T., Amuda, K., Olakunle, L., & Mercy, A.-A., T., 2022. The Challenges and Conservation Strategies of Biodiversity: The Role of Government and Non-Governmental Organization for Action and Results on the Ground, C. Izah S., (Eds.), *Biodiversity in Africa: Potentials, Threats, and Conservation*. Sustainable Development and Biodiversity. Springer. https://doi.org/10.1007/978-981-19-3326-4_18
- Raimi, M.O., Adeolu, A.T., Enabulele, C.E., Awogbami, S.O., 2018. Assessment of Air Quality Indices and its Health Impacts in Ilorin Metropolis. *Science Park Journals of Scientific Research and Impact*. 4(4), 60–74.
- Raimi, M.O., Bilewu, O.O., Adio, Z.O., Abdulrahman, H., 2019. Women Contributions to Sustainable Environments in Nigeria. *Journal of Scientific Research in Allied Sciences*. 5(4), 35–51. <https://doi.org/10.26838/JUSRES.2019.5.4.104>
- Raimi, M.O., Clinton, I.E., Olawale, H.S., 2021. Problematic Groundwater Contaminants: Impact of Surface and Ground Water Quality on the Environment in Ebocha-Obrikom Oil and Gas Producing Area of Rivers State, Nigeria, and others, (Eds.), 2nd International E-Conference on Geological and Environmental Sustainability, July 29-30, 2021. Oral Presentation Presented at the United Research Forum.
- Raimi, M.O., Ho, S., Isah, H.M., 2020. Health risk exposure to cypermethrin: A case study of kano state, *Journal of Agriculture*. 7th International Conference on Public Healthcare and Epidemiology.
- Raimi, M.O., Olawale, A.-A., Abiola, H.S., Abinotami, O.S., Ruth, W.E., Nimisingha, E.E., & Walter, D., O, B., 2022. Leaving No One Behind: Impact of Soil Pollution on Biodiversity in the Global South: A Global Call for Action, C. Izah S., (Eds.), *Biodiversity in Africa: Potentials, Threats and Conservation*. Sustainable Development and Biodiversity. Springer. https://doi.org/10.1007/978-981-19-3326-4_8
- Raimi, M.O., Raimi, A.G., 2020. The Toughest Triage in Decision Impacts: Rethinking Scientific Evidence for Environmental and Human Health Action in the Times of Concomitant Global Crises. *CPQ Medicine*. 11(1), 1–5.
- Raimi, M.O., Sawyerr, H.O., Ezekwe, I.C., Gabriel, S., 2022. Toxicants in Water: Hydrochemical Appraisal of Toxic Metals Concentration and Seasonal Variation in Drinking Water Quality in Oil and Gas Field Area of Rivers State, P.H. Saleh P.A.I. Hassan, (Eds.), *Heavy Metals - New Insights* [Working Title. <https://doi.org/10.5772/intechopen.102656>
- Raimi, M.O., Suleiman, R.M., Odipe, O.E., Salami, J.T., Oshatunberu, M., 2019. Women Role in Environmental Conservation and Development in Nigeria. *Ecology & Conservation Science*. 1(2), 1–16. <https://dx.doi.org/10.2139/ssrn.3425832>
- Raimi, O., Ezekwe, C., Bowale, A., Samson, T., 2022. Hydrogeochemical and Multivariate Statistical Techniques to Trace the Sources of Ground Water Contaminants and Affecting Factors of Groundwater Pollution in an Oil and Gas Producing Wetland in Rivers State,

- Nigeria. *Open Journal of Yangtze Oil and Gas*. 7, 166–202. <https://doi.org/10.1101/2021.12.26.21268415>
- Rosell, G., Quero, C., Coll, J., Guerrero, A., 2008. Biorational insecticides in pest management. *Journal of Pesticide Science*. 33, 103–121. <https://doi.org/10.1584/jpestics.R08-01>
- Samson, T.K., M, O.O., O, R., 2020. A Predictive Model for Confirmed Cases of COVID-19 in Nigeria. *European Journal of Applied Sciences*. 8(4), 1–10. <https://doi.org/10.14738/aivp.84.8705>
- Sosan, M.B., Oyekunle, J., Odewale, G., 2018. Occurrence and levels of organochlorine pesticide residues in maize samples from open markets and stores in Ile-Ife and Ondo, Southwestern Nigeria. *Nigeria Journal of Entomology*. 34(1), 25–37. [https://doi.org/10.36108/NJE/8102/43\(0140\)](https://doi.org/10.36108/NJE/8102/43(0140))
- Stathers, T.E., Arnold, S.E.J., Rumney, C.J., Hopson, C., 2020. Measuring the nutritional cost of insect infestation of stored maize and cowpea. *Food Security*. 12(2), 285–308. <https://doi.org/10.1007/s12571-019-00997-w>
- Stephen, O.A., Solomon, O.A., Henry, O.S., Afolabi, O.O., Morufu, O.R., 2022. Comprehensive Understanding of Hydrogeochemical evaluation of seasonal variability in groundwater quality Dynamics in the Gold Mining Areas of Osun State, Nigeria. medRxiv. <https://doi.org/10.1101/2022.11.06.22282015>
- Suleiman, R.M., Mo, R., Ho, S., 2019. A Deep Dive into the Review of National Environmental Standards and Regulations Enforcement Agency (NESREA) Act. *International Research Journal of Applied Sciences*, 2663–5585. [Irrjas.2019.123.123](https://doi.org/10.123.123)
- UNEP, 2008. Stockholm convention on persistent organic pollutants. United Nations Environment Programme (UNEP).
- Vanden Bilcke, C., 2002. The stockholm convention on persistent organic pollutants. *Review of European Comparative & International Environmental Law*. 11, 328–342.
- Zhang, Y.M., Xue, J., Zhai, J., Zhang, G.Q., Zhang, W.X., Wang, K.R., Ming, B., Hou, P., Xie, R.Z., Liu, C.W., Li, S.K., 2021. Does nitrogen application rate affect the moisture content of corn grains. *Journal of Integrative Agriculture*. 20(10), 63401–63404.